

# Analogy in mathematics

David Corfield

University of Kent

28 May, 2021

# Sir Michael Atiyah

*...mathematics can I think be viewed as the science of analogy.*


# Ulam reporting Banach

*A mathematician is a person who can find analogies between theorems; a better mathematician is one who can see analogies between proofs and the best mathematician can notice analogies between theories.*


# Ulam reporting Banach

*A mathematician is a person who can find analogies between theorems; a better mathematician is one who can see analogies between proofs and the best mathematician can notice analogies between theories. One can imagine that the ultimate mathematician is one who can see analogies between analogies.*

# Regular polygons



# Regular polyhedra




# Regular polytopes

2	regular polygons	triangle	square	...	...
3	5 Platonic solids	tetrahed.	cube	octahed.	2 more

# 4 dimensional polytopes

The 8-cell





# Regular polytopes

2	regular polygons	triangle	square	...	...
3	5 Platonic solids	tetrahed.	cube	octahed.	2 more
4	6 4-complexes	4-simplex	4-hypercube	4-orthoplex	3 more

# Regular polytopes

2	regular polygons	triangle	square	...	...
3	5 Platonic solids	tetrahed.	cube	octahed.	2 more
4	6 4-complexes	4-simplex	4-hypercube	4-orthoplex	3 more
5	3 5-complexes	5-simplex	5-hypercube	5-orthoplex	no more

n	simplex	hypercube	orthoplex
2	triangle	square	...
3	4-6-4	8-12-6	6-12-8
4	5-10-10-5	16-32-24-8	8-24-32-16
5	6-15-20-15-6	32-80-80-40-10	10-40-80-80-32

n	simplex	hypercube	orthoplex
2	3-3	4-4	4-4
3	4-6-4	8-12-6	6-12-8
4	5-10-10-5	16-32-24-8	8-24-32-16
5	6-15-20-15-6	32-80-80-40-10	10-40-80-80-32

2	regular polygons	triangle	square	...	$\infty$ more
3	5 Platonic solids	tetrahed.	cube	octahed.	2 more
4	6 4-complexes	4-simplex	4-hypercube	4-orthoplex	3 more
5	3 5-complexes	5-simplex	5-hypercube	5-orthoplex	no more
...	...	...	...	...	...
k	3 k-complexes	k-simplex	k-hypercube	k-orthoplex	no more

2	regular polygons	triangle	square	...	$\infty$ more
3	5 Platonic solids	tetrahed.	cube	octahed.	2 more
4	6 4-complexes	4-simplex	4-hypercube	4-orthoplex	3 more
5	3 5-complexes	5-simplex	5-hypercube	5-orthoplex	no more
...	...	...	...	...	...
k	3 k-complexes	k-simplex	k-hypercube	k-orthoplex	no more

Vaguer analogy: infinite families with sporadics, such as simple Lie groups ( $A_i, B_i, C_i, D_i, E_6, E_7, E_8, F_4, G_2$ ).

# Henri Poincaré

*...the mathematical facts worthy of being studied are those which, by their analogy with other facts, are capable of leading us to the knowledge of a mathematical law, just as experimental facts lead us to the knowledge of a physical law. They are those which reveal to us unexpected kinship between other facts, long known, but wrongly believed to be strangers to one another.*

# Henri Poincaré

*...the mathematical facts worthy of being studied are those which, by their analogy with other facts, are capable of leading us to the knowledge of a mathematical law, just as experimental facts lead us to the knowledge of a physical law. They are those which reveal to us unexpected kinship between other facts, long known, but wrongly believed to be strangers to one another.*

But we can have too much of a good thing...



# André Weil

*Nothing is more fruitful, every mathematician knows it, than these obscure analogies, these disputes reflected from one theory to another, these secret caresses, these quarrels for no reason; indeed nothing gives the researcher more pleasure. The day comes when the illusion falls away; intuition turns into certainty, twin theories reveal their common source, before vanishing; as the Gita tells us, we achieve knowledge and indifference at the same time. Metaphysics has become mathematics, fit content for a treatise whose austere beauty we no longer find moving.*

...

*Nothing is more fruitful, every mathematician knows it, than these obscure analogies, these disputes reflected from one theory to another, these secret caresses, these quarrels for no reason; indeed nothing gives the researcher more pleasure. The day comes when the illusion falls away; intuition turns into certainty, twin theories reveal their common source, before vanishing; as the Gita tells us, we achieve knowledge and indifference at the same time. Metaphysics has become mathematics, fit content for a treatise whose austere beauty we no longer find moving.*

...

*Fortunately for researchers, as soon as the mists dissolve in one place, they descend elsewhere.*

- Analogies indicating what may be expected to be subsumed under a covering law.

- Analogies indicating what may be expected to be subsumed under a covering law.
- Some are suggestive of a larger scheme, but resist complete systematisation (for now).

# Suggestive analogies, such as arithmetic topology

number ring $\text{Spec}(\mathcal{O}_k) \cup \{\infty\}$ $\text{Spec}(\mathbf{Z}) \cup \{\infty\}$	$\longleftrightarrow$	3-manifold $M$ $S^3$
prime : $\text{Spec}(\mathbf{F}_p) \subset \text{Spec}(\mathcal{O}_k)$ primes $p_1, \dots, p_n$ infinite prime	$\longleftrightarrow$	knot $K : S^1 \subset M$ link $K_1 \cup \dots \cup K_n$ end
$p$ -adic integers $\text{Spec}(\mathcal{O}_p)$	$\longleftrightarrow$	tube n.b.d $V(K)$
$p$ -adic field $\text{Spec}(k_p)$	$\longleftrightarrow$	torus $\partial V(K)$
$\pi_1(\text{Spec}(\mathcal{O}_p)) = \langle \sigma \rangle$	$\longleftrightarrow$	$\pi_1(V(K)) = \langle \beta \rangle$
$\pi_1^{\text{étale}}(\text{Spec}(k_p)) = \langle \tau, \sigma \mid \tau^{p-1}[\tau, \sigma] = 1 \rangle$	$\longleftrightarrow$	$\pi_1(\partial V(K)) = \langle \alpha, \beta \mid [\alpha, \beta] = 1 \rangle$



$k^{\times} \rightarrow \bigoplus_{p:\text{primes}} \mathbf{Z}$ $a \mapsto a \mathcal{O}_k^{\times}$	$\longleftrightarrow$	$C_2(M, \mathbf{Z}) \xrightarrow{\beta} C_1(M, \mathbf{Z})$ $\Sigma \mapsto \partial \Sigma$
class group $H_k$	$\longleftrightarrow$	$H_1(M, \mathbf{Z})$
$\mathcal{O}_k^{\times}$	$\longleftrightarrow$	$H_2(M, \mathbf{Z})$
$\pi_1(\text{Spec}(\mathcal{O}_k) \setminus \{p_1, \dots, p_n\})$ max. Galois group with given ramification	$\longleftrightarrow$	$\pi_1(M \setminus K_1 \cup \dots \cup K_n)$ link group
power residue symbol	$\longleftrightarrow$	linking number

This analogy likens the linking of loops in a 3-space with quadratic reciprocity.

This analogy likens the linking of loops in a 3-space with quadratic reciprocity.

The latter relates for two prime numbers,  $p$  and  $q$ , whether each is a square modulo the other.

This analogy likens the linking of loops in a 3-space with quadratic reciprocity.

The latter relates for two prime numbers,  $p$  and  $q$ , whether each is a square modulo the other.

There are suggestive 'dictionaries' for this analogy. But let's turn to the more precisely formulated end of analogy formation.



# Category theory as the great systematizer

Emily Riehl:

*Atiyah described mathematics as the “science of analogy.” In this vein, the purview of category theory is mathematical analogy. Category theory provides a cross-disciplinary language for mathematics designed to delineate general phenomena, which enables the transfer of ideas from one area of study to another. The category-theoretic perspective can function as a simplifying abstraction, isolating propositions that hold for formal reasons from those whose proofs require techniques particular to a given mathematical discipline.*

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
- Both are divisible by 12,

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
- Both are divisible by 12, but also by 9.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
- Both are divisible by 12, but also by 9.
- These latter both factor through 36.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
- Both are divisible by 12, but also by 9.
- These latter both factor through 36.
- All common divisors do this.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
- Both are divisible by 12, but also by 9.
- These latter both factor through 36.
- All common divisors do this.
  
- Consider two copies of the real numbers.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
  - Both are divisible by 12, but also by 9.
  - These latter both factor through 36.
  - All common divisors do this.
- 
- Consider two copies of the real numbers.
  - Consider a system with two real measurements such as the location and velocity of a pendulum,



# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
  - Both are divisible by 12, but also by 9.
  - These latter both factor through 36.
  - All common divisors do this.
- 
- Consider two copies of the real numbers.
  - Consider a system with two real measurements such as the location and velocity of a pendulum, but also the fertiliser load and crop yield of a collection of fields.

## GCD and the cartesian plane as 'products'

- Consider 72 and 108.
  - Both are divisible by 12, but also by 9.
  - These latter both factor through 36.
  - All common divisors do this.
- 
- Consider two copies of the real numbers.
  - Consider a system with two real measurements such as the location and velocity of a pendulum, but also the fertiliser load and crop yield of a collection of fields.
  - This data can be represented in the plane.

# GCD and the cartesian plane as 'products'

- Consider 72 and 108.
  - Both are divisible by 12, but also by 9.
  - These latter both factor through 36.
  - All common divisors do this.
- 
- Consider two copies of the real numbers.
  - Consider a system with two real measurements such as the location and velocity of a pendulum, but also the fertiliser load and crop yield of a collection of fields.
  - This data can be represented in the plane.
  - This is so for all such collections.

# Products

- Consider two propositions,  $P$  and  $Q$ .

# Products

- Consider two propositions,  $P$  and  $Q$ .
- Consider a proposition,  $R$ , that entails both of them.


# Products

- Consider two propositions,  $P$  and  $Q$ .
- Consider a proposition,  $R$ , that entails both of them.
- Then  $R$  entails  $P \wedge Q$ .

# Products

- Consider two propositions,  $P$  and  $Q$ .
- Consider a proposition,  $R$ , that entails both of them.
- Then  $R$  entails  $P \wedge Q$ .

All instances of:



# Preservation of products by right adjoints

- $\forall x(Px \& Qx)$  iff  $\forall xP(x) \& \forall xQ(x)$ .



# Preservation of products by right adjoints

- $\forall x(Px \& Qx)$  iff  $\forall xP(x) \& \forall xQ(x)$ .
- Given two polynomials, the zeros of  $f \times g = \text{zeros of } f + \text{zeros of } g$ .

# Preservation of products by right adjoints

- $\forall x(Px \& Qx)$  iff  $\forall xP(x) \& \forall xQ(x)$ .
- Given two polynomials, the zeros of  $f \times g =$  zeros of  $f +$  zeros of  $g$ .
- the set of points underlying the product of two spaces = the product of the sets of points underlying each space.

# Pólya's Bayesian account

- A is analogous to B
- A is true

# Pólya's Bayesian account

- A is analogous to B
- A is true
- Then B is somewhat more likely to be true

# Pólya's Bayesian account

- A is analogous to B
- A is true
- Then B is somewhat more likely to be true

Pólya speaks of the “hope for a common ground”,  $H$ .

# Pólya's Bayesian account

- A is analogous to B
- A is true
- Then B is somewhat more likely to be true

Pólya speaks of the “hope for a common ground”,  $H$ .

We face the standard problem of mere conjunction. We cannot simply have  $H = A \& B$ .

# Pólya's Bayesian account

- A is analogous to B
- A is true
- Then B is somewhat more likely to be true

Pólya speaks of the “hope for a common ground”,  $H$ .

We face the standard problem of mere conjunction. We cannot simply have  $H = A \& B$ .

$H$  must be ‘lawlike’.

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.



## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$
- $\sin(x)/x$  has roots  $x = \pm\pi, \pm2\pi, \pm3\pi, \dots$

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$
- $\sin(x)/x$  has roots  $x = \pm\pi, \pm2\pi, \pm3\pi, \dots$
- $\sin(x)/x = (1 - x^2/\pi^2)(1 - x^2/4\pi^2)(1 - x^2/9\pi^2)\dots?$

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$
- $\sin(x)/x$  has roots  $x = \pm\pi, \pm2\pi, \pm3\pi, \dots$
- $\sin(x)/x = (1 - x^2/\pi^2)(1 - x^2/4\pi^2)(1 - x^2/9\pi^2)\dots?$
- But then  $1 + 1/4 + 1/9 + \dots = \pi^2/6$ .

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$
- $\sin(x)/x$  has roots  $x = \pm\pi, \pm2\pi, \pm3\pi, \dots$
- $\sin(x)/x = (1 - x^2/\pi^2)(1 - x^2/4\pi^2)(1 - x^2/9\pi^2)\dots?$
- But then  $1 + 1/4 + 1/9 + \dots = \pi^2/6$ .

Euler checked this to many places. But it doesn't work for  $\tan(x)/x$ .

## Euler's analogy

We can relate the roots of an equation to the factorization of the relevant polynomial.

- If  $f(x) = 0$  has roots  $\alpha$  and  $\beta$ , then  $f(x) = A(1 - x/\alpha)(1 - x/\beta)$ .
- Works for any finite number of roots.
- Could this work for infinitely many roots?
- $\sin(x)/x = 1 - x^2/3! + x^4/5! - \dots$
- $\sin(x)/x$  has roots  $x = \pm\pi, \pm2\pi, \pm3\pi, \dots$
- $\sin(x)/x = (1 - x^2/\pi^2)(1 - x^2/4\pi^2)(1 - x^2/9\pi^2)\dots?$
- But then  $1 + 1/4 + 1/9 + \dots = \pi^2/6$ .

Euler checked this to many places. But it doesn't work for  $\tan(x)/x$ .

Eventually a general theorem (a common ground) concerning such factorisation appears in the 19th century.



# Riemann hypothesis

The real part of every non-trivial zero of the Riemann zeta function is  $\frac{1}{2}$ .

# Riemann hypothesis

The real part of every non-trivial zero of the Riemann zeta function is  $\frac{1}{2}$ .

- Checked by computer for billions of zeros.
- An analogue (Rosetta Stone) for finite fields is known to be true.

# Riemann hypothesis

The real part of every non-trivial zero of the Riemann zeta function is  $\frac{1}{2}$ .

- Checked by computer for billions of zeros.
- An analogue (Rosetta Stone) for finite fields is known to be true.

*There is impressive numerical evidence in its favour but certainly the best reason to believe that it is true comes from the analogy of number fields with function fields of curves over finite fields where the analogue of RC has first been proved by A. Weil. (Deninger 1994)*

[Developed further by Deligne.]

## Caution needed, Munson

*We intend to explain some of the intuition behind one incarnation of calculus of functors, namely the so-called “manifold calculus” due to Weiss and Goodwillie. Specifically, we will **highlight some analogies** between the ordinary calculus of functions and the manifold calculus of functors. The **trouble with analogies is that they are not equivalences**, and some may lead the reader to want to push them further. Some may indeed be pushed further than we are currently aware, and some may lead to direct contradictions and/or bad intuition. Another risk is that it is considered bad manners to tell people how to categorize various ideas: part of our mathematical culture seems to be that we leave intuition for talks and personal communications and rigor and precision for our papers, and with good reason: we cannot anticipate the ways in which our work may be useful in the future, and so it may be best to convey it in as concise and precise a way as possible. We feel the relatively small risk of misleading the reader and the faux pas of making permanent intuitive notions by publishing them is a small price to pay for the possibility that this may entice some reader to learn more about these ideas and try to use them. The structure of the category...is much richer than the usual topology on the real line, so **analogies between functions and functors may seem a little weak**. Still, there are a few things to say that may be helpful. We like to think of ... as the **analog** of the dense subset  $Q$ .*

## To conclude

- Analogical reasoning is everywhere in maths.
- Category theory is good at capturing very general, precise analogies across fields.
- Some analogies run for a very long time without being turned into general theory.
- Analogies play a huge role in plausibility assessment.
- Care must be taken not to follow them blindly.