



# Introduction to Quark Gluon Plasma

---

Mahnaz Q. Haseeb  
Department of Physics  
CIIT, Islamabad



# Topics of lectures on QGP

---

- Introduction to QGP
- Thermodynamics and Phase Transitions
- Hydrodynamics in QGP



# Outline


---

- Motivation
- An overview of QCD
- What is Quark Gluon Plasma (QGP)?
- How to create QGP?
- Claims about its discovery
- Physical parameters measured
- Expectation from LHC about QGP

# Motivation

“To understand the Equation of State of Nuclear, Hadronic & Partonic Matter”

Of cross-disciplinary interest:



## Nuclear Physics “NMEOS”

Collective Nuclear Phenomena  
Effects of the Nuclear Medium

## Astrophysics “NMEOS - Matter Incompressibility”

Neutron Star Stability  
Supernova Expansion Dynamics



## Cosmology “QCD Phase Transition”

Evolution of Early Universe

## Particle Physics “Perturbative QCD Vacuum”

High (Energy) Density QCD  
Symmetry Breaking Mechanisms  
Particle Masses

# History of the Universe



Source: Nuclear Science Wall Chart

# Energy Scales

---

- **The beginning**

The universe is a hot plasma of fundamental particles ... quarks, leptons, force mediating particles (and other particles ?)

$10^{-43}$  s      Planck scale (quantum gravity ?)       $10^{19}$  GeV

$10^{-35}$  s      Grand unification scale (strong+electroweak)       $10^{15}$  GeV

Inflationary period  $10^{-35}$ - $10^{-33}$  s

$10^{-11}$  s      Electroweak unification scale      200 GeV

- **Micro-structure**

$10^{-5}$  s      QCD scale - protons and neutrons form      200 MeV

3 mins      Primordial nucleosynthesis      5 MeV

$3 \times 10^5$  y      Radiation and matter decouple - atoms form      1 eV

- **Large scale structure**


1 b yrs      Proto-galaxies and the first stars


3 b yrs      Quasars and galaxy spheroids

5 b yrs      Galaxy disks

Today      Life !

# A brief history ...





$\approx 1$  cm  
**Crystal**

Size Ratio:  $1/10,000,000$

$10^{-7}$  cm  
**Molecule** ←

$1/10$

$10^{-8}$  cm  
**Atom**

$1/10,000$

$10^{-13}$  cm  
**Atomic nucleus**


$1/10$

$10^{-13}$  cm  
**Proton**

$1/1,000$

$< 10^{-16}$  cm  
**Electron,  
Quark** →

# Building Blocks of Hadron World



Strong interaction is due to color charges and mediated by gluons. Gluons carry color charges too.


Baryon Density:  $\rho = \text{baryon number/volume}$

normal nucleus  $\rho_0 \sim 0.15 \text{ /fm}^3 \sim 0.25 \times 10^{15} \text{ g/cm}^3$

Temperature:  $\text{MeV} \sim 1.16 \times 10^{10} \text{ K}$

$10^{-6}$  second after the Big Bang  $T \sim 200 \text{ MeV}$

# Interaction between quarks and gluons




Separate quarks do not exist in nature

- confinement first established as an experimental fact, then by theory of strong interactions

**Quantum Chromo Dynamics (QCD)**

# Analogies and differences between QED and QCD

to study structure of an atom...




neutral atom

...separate constituents


QED Quantum Electro Dynamics

**Confinement:** fundamental & crucial (but *not* understood!) feature of strong force  
- colored objects (quarks) have  $\infty$  energy in normal vacuum



QCD Quantum Chromo Dynamics

# QCD Potential




## in vacuum:

- linear increase with distance, strong attractive force
- confinement of quarks to hadrons

## in dense and hot matter

- screening of color charges (similar to Debye screening in dense atomic matter)
- potential vanishes for large distance
- deconfinement of quarks  
→ QGP

- 
- An electrical plasma is one in which the particles are charged
    - a nuclear plasma would be one in which the particles possess color charges.
  - The charge of one particle is screened by the surrounding charges.
  - **Debye Screening Radius ( $\lambda_D$ ):**  
The distance at which the charge is reduced by  $1/e$  for electromagnetic plasma.



These quarks effectively cannot “see” each other!

# Instances of Debye Screening

---

## ○ **In bulk media**

- In bulk media, there is an additional charge screening effect.
- At high charge density,  $n$ , the short range part of the potential becomes:

$$V(r) \propto \frac{1}{r} \Rightarrow \frac{1}{r} \exp\left[\frac{-r}{r_D}\right] \quad \text{where} \quad r_D = \frac{1}{\sqrt[3]{n}}$$

and  $r_D$  is the Debye screening radius.

- Effectively, long range interactions ( $r > r_D$ ) are screened.

## ○ **The Mott transition**

- In condensed matter, when  $r <$  electron binding radius  
⇒ an electric insulator becomes conducting.


## ○ **Debye screening in QCD**

- Analogously, think of the quark-gluon plasma as a colour conductor.
- Nucleons (all hadrons) are colour singlets (baryons:  $qqq$ , or mesons  $q\bar{q}$ ).
- At high (charge) density quarks and gluons become unbound.  
⇒ nucleons (hadrons) cease to exist.


# Debye screening

- Modification of  $V_{em}$  - the Mott Transition







a)  $d > r_D$



b)  $d < r_D$




# Screening in QCD



# What is Quark-Gluon Plasma?

At room temperature, quarks and gluons are always confined inside **colorless** objects (**hadrons**):

protons, neutrons, pions, .....




Very high temperature (asymptotic freedom):

- Interactions become weak
- quarks and gluons **deconfined**
- **Quark-gluon plasma (QGP)**

Infinitely high temperature:

QGP may behave like an **ideal gas**.




# Generating a deconfined state

---

Present understanding of Quantum Chromodynamics (QCD)

- heating
  - compression
- *deconfined matter !*





Nuclear Matter  
(confined)

Hadronic Matter  
(confined)

Quark Gluon Plasma  
deconfined !


# Where to study the QGP?



# Relation to the Big Bang


- Energy density
  - 1 GeV/fm<sup>3</sup> →
  - 1.8x10<sup>15</sup>g/cm<sup>3</sup>
- Temperature
  - 170 MeV →
  - 2.0x10<sup>12</sup> K

Conditions that prevailed ≈ 10 μs after the Big Bang




# The Melting of Quarks and Gluons

**Matter Compression:**



High Baryon Density  
-- low energy heavy ion collisions  
-- neutron star  $\rightarrow$  quark star






**Vacuum Heating:**



High Temperature Vacuum  
-- high energy heavy ion collisions  
-- the Big Bang

**Deconfinement**


# Creation of QGP




**Collide ultra-relativistic heavy ions**

Pb+Pb 160 GeV/A

$t = -17.33$  fm/c





UrQMD Frankfurt/M












Pb+Pb 160 GeV/A


$t = 24.17$  fm/c



UrQMD Frankfurt/M

Pb+Pb 160 GeV/A





$t = 43.17 \text{ fm}/c$




UrQMD Frankfurt/M

# The "Little Bang" in the lab

- *Nucleus-nucleus collisions*: fixed-target reactions ( $\sqrt{s}=20$  GeV, SPS) or colliders ( $\sqrt{s}=200$  GeV, RHIC.  $\sqrt{s}=5.5$  TeV, LHC)
- *QGP* expected to be formed in a *tiny region* ( $\sim 10^{-14}$  m) and to last *very short times* ( $\sim 10^{-23}$  s).
- *Collision dynamics*: Diff. observables sensitive to diff. react. stages





# Time in Heavy Ion Reactions




## **p+p interactions:**

- No extended initial medium
- Chemical freeze-out (no thermalization)
- Kinetic freeze-out close to chemical freeze-out

Particle yields


Particle spectra



## **Au+Au interactions:**


- Extended hot and dense phase
- Thermalization at chem. freeze-out
- Kinetic freeze-out separated from chemical freeze-out

# Ultrarelativistic Heavy Ion Collisions



# End of the World!

J. Nagle



Can be dismissed with some basic General Relativity

$$R_S = \frac{2GM}{c^2} = 10^{-49} \text{ meters}$$

$$R = 10^{-15} \text{ meters}$$

much less  
than  
Planck length  
!

Even if it could form, it would evaporate by Hawking Radiation in  $10^{-83}$  seconds !

# A little of search history

---

**1951-1975**

**Pomeranchuk, Hagedorn,  
Fubini, Veneziano,  
Mandelstam, Cabibbo, Parisi  
et al:**

**Critical temperature for  
phase transition to QGP  
with free color quarks and  
gluons**

$$T_c = 150-180 \text{ MeV}$$

**1986/1987**

**Experimental research of AA  
interactions:**

**Creation of new state of matter  
– Quark Gluon Plasma at AGS  
and SPS, since 2000 at RHIC  
and now shortly in LHC**


**Density of energy in point of  
transition**

$$0.7-1.0 \text{ GeV/fm}^3$$

$$\text{AGS} : \sqrt{S_{NN}} = 5 \text{ GeV}$$

$$\text{SPS} : \sqrt{S_{NN}} = 20 \text{ GeV}$$

$$\text{RHIC} : \sqrt{S_{NN}} = 200 \text{ GeV}$$



# Facilities to hunt the QGP and its signatures

-What were the expectations in the beginning?

Little Bang

The Future Experiment

CBM: Fixed Target  
 $U^{235}$  @ 2-45 GeV/A

2009-10

LHC: Collider  
Pb+Pb @ 5.5 TeV/A

2000

RHIC: Collider  
Au+Au @ 200 GeV/A

1996

SPS: Fixed Target  
Pb at 158 GeV/A  
(Ec.m.=17.3 GeV)

1991

AGS: Fixed Target  
Au at 11.7 GeV/A  
(Ec.m.=4.86 GeV)

1984


Bevalac:  
Fixed Target  
Au at  
1 GeV/A

- What have we learned so far?
- What are the prospects for the future?

# QGP: A new state of matter

CERN  
Organisation Européenne pour la Recherche Nucléaire  
European Organization for Nuclear Research

**New State of Matter created at CERN**



PRESS RELEASE

“The combined data coming from the seven experiments on CERN's Heavy Ion programme have given a clear picture of a new state of matter. . . . We now have evidence of a new state of matter where quarks and gluons are not confined. There is still an entirely new territory to be explored concerning the physical properties of quark-gluon matter.” [ L. Maiani, 2000]

# Research results, SPS:

**February 2000**

**Ulrich Heinz and Maurice Jacob Theoretical Physics Division, CERN CH-1211 Geneva 23, Switzerland:**

**Evidence for a New State of Matter:**

---

**An Assessment of the Results from the CERN Lead Beam**

... Present theoretical ideas provide a more precise picture for this new state of matter: it should be **a quark-gluon plasma (QGP), in which quark and gluons, the fundamental constituents, are no longer confined within the dimensions of the nucleon, but free to move around over a volume in which a high enough temperature and/or density prevails...**

... A common assessment of the collected data leads us to conclude that we now have compelling evidence that a new state of matter has indeed been created... **The new state of matter found in heavy ion collision at the SPS features many of the characteristics of the theoretically predicted quark-gluon plasma...**

# For last 9 years there has been a significantly increased interest in HIC

---

- New **beautiful results from RHIC** (STAR, PHENIX, BRAHMS, PHOBOS), and also **new results from experiments at SPS** (NA49, NA50, NA57 and NA60).
- **Ability to deliver physics at ~ all scales**

**Research program on nucleus-nucleus collisions at the new accelerator facility at GSI (FAIR) and LHC**

- **There is very fast development of theory and preparation of new experiments**

# SCIENTIFIC AMERICAN

MAY 2006  
WWW.SCIAM.COM

## Quark Soup

PHYSICISTS RE-CREATE  
THE LIQUID STUFF OF  
**THE EARLIEST  
UNIVERSE**



search BNL

Go

Newsroom Links

- Newsroom Home
- Search News Archives
- Photo Archive
- Streaming Video
- Fact Sheets
- Science Magazine
- Management Bios
- About Brookhaven

Contact: Karen McNulty Walsh, (631) 344-8350 or Mona S. Rowe, (631) 344-5056

Print-friendly E-mail Article

## RHIC Scientists Serve Up "Perfect" Liquid

### New state of matter more remarkable than predicted -- raising many new questions

April 18, 2005

TAMPA, FL -- The four detector groups conducting research at the [Relativistic Heavy Ion Collider](#) (RHIC) -- a giant atom "smasher" located at the U.S. Department of Energy's Brookhaven National Laboratory -- say they've created a new state of hot, dense matter out of the quarks and gluons that are the basic particles of atomic nuclei, but it is a state quite different and even more remarkable than had been predicted. In [peer-reviewed papers](#) summarizing the first three years of RHIC findings, the scientists say that instead of behaving like a gas of free quarks and gluons, as was expected, the matter created in RHIC's heavy ion collisions appears to be more like a *liquid*.

"Once again, the physics research sponsored by the Department of Energy is producing historic results," said Secretary of Energy Samuel Bodman, a trained chemical engineer. "The DOE is the principal federal funder of basic research in the physical sciences, including nuclear and high-energy physics. With today's announcement we see that investment paying off."



"The truly stunning finding at RHIC that the new state of matter created in the collisions of gold ions is more like a liquid than a gas gives us a profound insight into the earliest moments of the universe," said Dr. Raymond L. Orbach, Director of the DOE Office of Science.

XML Get BNL News via RSS


# Key Results from SPS & RHIC


---

- **SPS** (1986-2003) – QGP signatures seen e.g.
  - **Enhancement of strangeness** (up to a factor of  $\sim 20$ ).
  - **Suppression of  $J/\Psi$  (c c-bar meson)**.
- **RHIC** (2000 onwards) – results from SPS exps confirmed, plus
  - **Elliptic flow**: collective flow of final state hadrons w.r.t. the reaction plane – medium behaves like an ideal liquid rather than a gas as expected.
  - **Jet quenching**: suppression of high  $p_T$  hadrons w.r.t. yield expected from superposition of nucleon-nucleon collisions.

- 
- Charmonium states are screened in QGP
    - $J/\psi$  suppression

### $J/\psi$ Suppression in QGP





# Large quantitative gains

---

Increasing the center of mass energy implies

- ➔ Denser initial system
- ➔ Longer lifetime
- ➔ Bigger spatial extension
- ➔ Stronger collective phenomena

A large body of experimental data from the CERN SPS and RHIC supports this argument.

# ALICE - A Large Ion Colliding Experiment

---

- LHC is **ultimate machine** for Heavy Ion Collisions
  - very **significant step** beyond RHIC
  - excellent conditions for **experiment & theory** (QCD)
  - not only latest, but **possibly last** HIC setup at the energy frontier
- ALICE is a powerful **next generation detector**
  - first truly **general purpose** Heavy Ion experiment
  - many **evolutionary** developments
    - SSD, SDD, TPC, em cal, ...
  - some **big advances** in technology
    - electronics, pixels, TOF, computing

# Pb+Pb Collisions at LHC

---

- cm-energy  $\sqrt{s} = 1150 \text{ TeV} = 0.18 \text{ mJ}$
- energy density  $\varepsilon = 1000 \text{ GeV/fm}^3$   
( $\varepsilon_{\text{Pb}} = 0.15 \text{ GeV/fm}^3$ )
- initial temperature  $T = 1 \text{ GeV}$   
( $T_{\text{critical}} = 0.15 \text{ GeV}$ )
- total multiplicity = 60000
- total volume at freeze-out  $V = 6 \cdot 10^5 \text{ fm}^3$   
( $V_{\text{Pb}} = 1500 \text{ fm}^3$ )
- lifetime until freeze-out  $\tau = 50 \text{ fm}/c$

# ALICE Program and Outlook

---

Safarik, 09

- first pp run
  - **important pp reference data for heavy ions**
  - **unique physics to ALICE**
    - **minimum-bias running**
    - **fragmentation studies**
    - **baryon-number transport**
    - **heavy-flavour cross sections**
- first few heavy-ion collisions
  - **establish global event characteristics**
  - **important bulk properties**
- first long heavy-ion run
  - **quarkonia measurements**
  - **Jet-suppression studies**
  - **flavour dependences**

## Outlook

- high luminosity heavy ion running ( $1\text{nb}^{-1}$ )
  - **dedicated high  $p_t$  electron triggers**
  - **jets > 100 GeV (EMCAL)**
  - **Y - states**
  - **$\gamma$  - jet correlations**
  - ...
- pA & light ion running

# ALICE @ LHC

---

- Evidence for QGP formation at CERN SPS and RHIC energies.
- ALICE will be able to study the physics of quark matter in detail.
  - almost all known observables
  - from early to late stages of QGP
- ALICE to study pp physics in its own right
- ALICE is ready for first physics


----- looking forward to lots of exciting physics shortly

# Jets – a new observable at LHC

- Hard, perturbative scale:  $Q \gg \Lambda_{QCD}$ .
- Parton shower development affected by the medium
- At LHC in Pb+Pb collisions:
  - ▶ wider  $p_T$  range for suppression, quenching studies
  - ▶ jet structure will likely be modified compared to jets in p+p
    - ▶ comparison to p+p and p+A is essential

## ➤ Observables:

- ▶ High  $p_T$  particles and particle correlations
- ▶ Jet rates: single and multi-jets (quenching studies)
- ▶ Jet fragmentation and shape:
  - ▶ Distance  $R$  to leading particle (in  $\eta$ - $\phi$  space)
  - ▶ forward-backward correlation:  $\Delta\phi$  (particle, jet axis)
  - ▶ Fragmentation function:  $F(z) = 1/N_j \times dN_{ch}/dz$  where  $z = p_t/p_{jet}$
- ▶ correlations with non-hadronic particles: jets+ $\gamma$ , jets+Z
- ▶ Jets originating from heavy quarks (b, c)




# EVIDENCE FOR A DENSE LIQUID (from Riordan and Zajc, Scientific American, May 2006, p. 34a.)


Two phenomena in particular point to the quark-gluon medium being a dense liquid state of matter: jet quenching and elliptic flow. Jet quenching implies the quarks and gluons are closely packed, and elliptic flow would not occur if the medium were a gas.

## ELLIPTIC FLOW

Off-center collisions between gold nuclei produce an elliptical region of quark-gluon medium.




The pressure gradients in the elliptical region cause it to explode outward, mostly in the plane of the collision (arrows).




## JET QUENCHING

In a collision of protons, hard scattering of two quarks produces back-to-back jets of particles.



In the dense quark-gluon medium, the jets are quenched, like bullets fired into water, and on average only single jets emerge.





---

Finally, lots of exciting and challenging physics expected from LHC experiments ...  
Best summarised by the following:

**“Traveller, road is nothing more than your footprints;  
Traveller, there is no road, you make it as you go”.**

H. Satz (*hep-ph/0209181*)



Thankyou

# Heavy-ion physics with ALICE

- **fully commissioned detector & trigger**
  - alignment, calibration available (pp)
- **first  $10^5$  events:** global event properties
  - multiplicity, rapidity density
  - elliptic flow
- **first  $10^6$  events:** source characteristics
  - particle spectra, resonances
  - differential flow analysis
  - interferometry
- **first  $10^7$  events:** high- $p_t$ , heavy flavours
  - jet quenching, heavy-flavour energy loss
  - charmonium production
- **yield bulk properties of created medium**
  - energy density, temperature, pressure
  - heat capacity/entropy, viscosity, sound velocity, opacity
  - susceptibilities, order of phase transition
- **early ion scheme**
  - 1/20 of nominal luminosity
  - $\int L dt = 5 \cdot 10^{25} \text{ cm}^{-2} \text{ s}^{-1} \times 10^6 \text{ s}$   
0.05 nb $^{-1}$  for PbPb at 5.5 TeV
  - $N_{pp \text{ collisions}} = 2 \cdot 10^8$  collisions  
400 Hz minimum-bias rate  
20 Hz central (5%)
  - muon triggers:  
 $\sim 100\%$  efficiency, < 1kHz
  - centrality triggers:  
bandwidth limited  
 $N_{PbPb \text{ minb}} = 10^7$  events (10Hz)  
 $N_{PbPb \text{ central}} = 10^7$  events (10Hz)

# Status: Was The QGP Seen?

---

- observations at SPS
  - initial energy density above critical value
  - strong collective effects
    - strongly interacting system
  - strangeness enhancement
    - close to equilibrium - possible in hadronic system?
  - photon radiation?
    - hot system in initial state - nature of system not conclusive

- 
- 
- modified vector meson spectra
    - relation to chiral symmetry restoration?
  - $J/\psi$ - suppression
    - currently better explained by QGP - unique?
  - no jet quenching
    - no strong effect expected

- 
- 
- observations at RHIC
    - initial energy density high above critical value
  - very strong collective effects
    - early equilibration
  - strangeness enhancement
    - as at SPS
  - photon radiation not yet observed
  - vector meson spectra not yet observed
  - $J/\psi$  - suppression not yet observed
  - jet quenching
    - high color charged density - nature of system not conclusive

# ALICE Summary & Outlook

---

first pp run

- **important pp reference data for heavy ions**
- **unique physics to ALICE**
  - **minimum-bias running**
  - **fragmentation studies**
  - **baryon-number transport**
  - **heavy-flavour cross sections**
- first few heavy-ion collisions
  - **establish global event characteristics**
  - **important bulk properties**
- first long heavy-ion run
  - **quarkonia measurements**
  - **Jet-suppression studies**
  - **flavour dependences**

## Outlook

- high luminosity heavy ion running ( $1\text{nb}^{-1}$ )
  - **dedicated high  $p_t$  electron triggers**
  - **jets  $> 100$  GeV (EMCAL)**
  - **$Y$  - states**
  - **$\gamma$  - jet correlations**
  - ...
- pA & light ion running

# Why Heavy Ions?

- Higher energy density may be achieved in proton-proton, but the partonic re-interaction time scale of order 1 fm/c.
- It is difficult to select events with different geometries and avoid autocorrelations.
- We will see that probes with long paths through the medium are key.

We should not rule out pp reactions, but rather study the similarities and differences with AA reactions.

# @ SPS

- All indications are that deconfinement is seen @SPS
- strangeness enhancement and  $J/\Psi$  suppression are correlated ( $\gamma_S$  vs centrality) !!
- SPS offers the unique possibility to study precisely the onset of deconfinement....
- to be considered in long term planning of the SPS!

# @ RHIC

- Deconfined phase is showing unexpected properties
- New phenomena, new probes:
  - jet tomography
  - collective motion
  - b-quarkonia could be a useful probe
- Initial quanta: Color Glass Condensate?
- A very dense, fluid phase: strongly interacting Quarks and Gluons?
- Which excitations populate the QG Liquid?

- RAPIDITY.

The rapidity is a generalization of the velocity.

$$y \equiv \operatorname{arcth}(v_{\parallel}) = \operatorname{arcth}\left(\frac{p_{\parallel}}{p^0}\right) = \frac{1}{2} \ln\left(\frac{p^0 + p_{\parallel}}{p^0 - p_{\parallel}}\right) \quad (2.5)$$

This definition uses the components of vectors  $\vec{v}$  and  $\vec{p}$

### 2.1.2 Properties of the rapidity

For small velocities:  $y \approx v_{\parallel}$ . Particle after the collision moves with  $\vec{v}$ .

It is customary to decompose  $\vec{v}$  to coordinates  $(y, \vec{p}_{\perp}/m)$ .

$\vec{p}$  can also be decomposed as  $p^{\mu} = (p^0, p_{\parallel}, \vec{p}_{\perp})$ .

The limit of rapidity coordinates for non-relativistic velocities:  $(y, \vec{p}_{\perp}/m) \rightarrow (v_{\parallel}, \vec{v}_{\perp})$ .

While the velocity is limited to 1 ( $c$ ), the rapidity  $y$  may vary between  $(-\infty, \infty)$ . See Fig. 2.4.




Figure 2.4 *The beam-parallel component of the velocity as function of the rapidity  $y$ , [2]*



---

## Review of last lecture: Photons

---

- A hot QGP will emit photons
- Once emitted, photons leave system
- But any hot system, QGP or hadrons, will emit photons
  - if contained in box, cannot use photon spectrum to distinguish QGP vs. hadrons
  - if  $T_{\text{photon}} > 200 \text{ MeV}$  unlikely to be from hadrons
  - closer analogy is box with transparent walls
    - photons not in thermal equilibrium
- photons extremely difficult to measure
  - large background of e.g.  $\pi^0 \rightarrow 2\gamma$



---

## Today

---

- High  $p_t$  signals of QGP
  - energy loss of partons within a plasma
- Schedule
  - Nov 11  $J/\psi$
  - Nov 18 strangeness
  - Nov 25 Thanksgiving - no class
  - Dec 2 HBT
  - Dec 9 low- $p_t$  and DCC



---

## Summary

---

- High pt partons lose energy in a plasma
  - scattering and produced gluons
  - amplitudes interfere (LPM effect)
    - lowers e-loss,  $dE/dx \sim \text{length}$
- Softens hadron spectra from fragmented parton
  - measure centrality dependence, periph. baseline
  - uncertainties, establish from p+A
    - pre-scattering: Cronin effect
    - gluon structure functions in Au: initial state
- Sensitive to density of colored-scattering centers

## Summary

---

- Charmonium states screened in a QGP  $\Rightarrow$  suppression
- pp reactions produce pre-resonant, color octet state  $\Rightarrow$  evolves into  $J/\psi$ ,  $\chi$  etc.
- Explains pp, pA and A+A data for  $A \leq$  sulphur
- NA50 Pb+Pb data has fewer  $J/\psi$ 
  - smooth dependence with  $E_t$ , no discontinuity
  - hadronic model reproduces data
    - early role of heavy hadrons, formation time of hadrons?
  - QGP ‘model’ reproduces data
    - can plasma be opaque to  $J/\psi$ , no e-loss of hard partons?
  - measure pt spectra of  $J/\psi$



---

## Summary

---

- Strangeness enhancement largest at lower  $E_{\text{beam}}$ 
  - hadronic mechanisms dominate
- Any QGP signal would sit on this baseline
  - strange-yields not reproduced by hadronic transport models
  - but without a definitive ‘smoking gun’, how bad a failure before you ‘need’ a QGP
  - key is excitation function 1-10, 40, 160 AGeV
    - ideal would be a minimum in enhancement
    - imply a QGP-driven enhancement above minimum



---

## Summary

---

- low-pt sensitive to large-scale phenomena
  - emission from cold QGP
  - collective motion in plasma
  - decay of disoriented vacuum
- any scattering after hadronization will remove info.
- low-pt in pA, AA, consistent decay of resonances
  - difference  $\pi^-$ ,  $\pi^+$  Coulomb interaction  $\Rightarrow$  size
- experimental difficulties
- RHIC: opportunities across several experiments

# Statistical QCD

$$\varepsilon_{QED} = \frac{\pi^2}{30} \left[ 2 + \frac{7}{8} 2 \times 2 \right] T^4$$

photon spin

electrons spin

$$\varepsilon_{QCD} = \frac{\pi^2}{30} \left[ 3 \times 8 + \frac{7}{8} \times 2 \times 2 \times 3 \times 3 \right] T^4$$

gluon spin, color

quarks spin, color, flavor

**Energy density reflects the information on what the matter is made of !**