

ON THE BURNSIDE RING AND STABLE COHOMOTOPY OF A FINITE GROUP

ERKKI LAITINEN*

0. Introduction.

In this paper we study the connection between permutation representations of finite groups G and stable cohomotopy of the classifying space BG , analogous to the connection between the character ring of G and the K -theory of BG .

Suppose S is a finite G -set. Each ordering of S gives a homomorphism $\varrho: G \rightarrow \Sigma_{|S|}$, where Σ_n denotes the permutation group in n letters and $|S|$ is the cardinality of S . Different orderings give conjugate maps, as do isomorphic G -sets. Hence the homotopy class of $B\varrho: BG \rightarrow B\Sigma_{|S|}$ only depends on the isomorphism class of S .

The disjoint union $\coprod_{n \geq 0} B\Sigma_n$ is a monoid and its group completion $\Omega B(\coprod_{n \geq 0} B\Sigma_n)$ is homotopy equivalent (as an H -space) to the space $QS^0 = \lim_{n \rightarrow \infty} \Omega^n S^n$ of stable self maps of spheres. Let $i: \coprod_{n \geq 0} B\Sigma_n \rightarrow QS^0$ be the resulting H -map and form the composition

$$\alpha_G(S): BG \rightarrow B\Sigma_{|S|} \rightarrow \coprod_{n \geq 0} B\Sigma_n \xrightarrow{i} QS^0.$$

Disjoint union and Cartesian product turn the equivalence classes of G -sets into a semiring, whose associated ring is the Burnside ring $A(G)$ ([17], [7]), and the correspondence $S \rightarrow \alpha_G(S)$ defines an additive map

$$\alpha_G: A(G) \rightarrow [BG, QS^0].$$

$[BG, QS^0]$ is by definition the stable cohomotopy $\pi_S^0(BG)$.

The space QS^0 admits besides the loop addition the smash product, homotopic to the product given by composition of maps. If $\coprod_{n \geq 0} B\Sigma_n$ is equipped with the monoid structure induced from the homomorphisms $\Sigma_n \times \Sigma_m \rightarrow \Sigma_{nm}$, then i respects both structures and α_G is a ring homomorphism.

The space QS^0 splits into a disjoint union of homotopy equivalent spaces $Q_n S^0$, $n \in \mathbb{Z}$, where $Q_n S^0$ denotes the subspace of degree n maps. Thus we have

* Partially supported by "Nordisk Universitetsstipendium" at Aarhus University.

Received March 3, 1978.

an augmentation $[BG, QS^0] \rightarrow \mathbf{Z}$. Taking cardinality of G -sets defines an augmentation of $A(G)$, and α_G is clearly augmentation preserving. The $(n+1)$ -fold products become trivial on the n -skeleton $B_n G$, so α_G factors as

$$\alpha_G: A(G)/I^{n+1}(G) \rightarrow [B_n G, QS^0].$$

Passing to the limit we get a map from the $I(G)$ -adic completion

$$\hat{\alpha}_G: \hat{A}(G) \rightarrow \varprojlim [B_n G, QS^0] = [BG, QS^0]$$

(the last isomorphism follows from the finiteness of $[B_n G, Q_0 S^0]$).

The map $\hat{\alpha}_G$ is analogous to the isomorphism from the completed representation ring $\hat{R}(G)$ to $K(BG)$ [1], and some time ago G. Segal made the

CONJECTURE. $\hat{\alpha}_G: \hat{A}(G) \rightarrow [BG, QS^0]$ is an isomorphism.

The full conjecture seems very hard and is probably out of reach at the moment. In this paper we study the injectivity of $\hat{\alpha}_G$.

First we reduce the problem to p -groups by showing

THEOREM A. *If $\hat{\alpha}_{G_p}$ is injective for the Sylow subgroups G_p of G , then $\hat{\alpha}_G$ is injective.*

This is proved by showing that $\hat{A}(G)$ embeds into $\bigoplus_p \hat{A}(G_p)$ via the restriction maps, and compatibly with $BG_p \rightarrow BG$.

For cyclic groups the natural map $\hat{A}(G) \rightarrow \hat{R}(G)$ is injective, and using Atiyah's result, $\hat{R}(G) \cong K^*(BG)$, we deduce

THEOREM B. $\hat{\alpha}_G$ is injective for cyclic groups G .

One cannot hope to detect the maps $\alpha_G(x): BG \rightarrow Q_0 S^0$ for $x \in \text{Ker}(\hat{A}(G) \rightarrow \hat{R}(G))$ by K -theory, since the space $Q_0 S^0$ splits as $J \times \text{cok } J$ with $\text{cok } J$ a K -theory point, and at least for groups G of odd order $[BG, J]$ embeds into $[BG, BU \times \mathbf{Z}] = \hat{R}(G)$. By studying induced maps in homology we prove

THEOREM C. $\hat{\alpha}_G$ is injective for elementary abelian groups $G = (\mathbf{Z}/p)^n$.

For Theorem C we need an induction machine, which tells that if α_H is injective for all genuine subgroups H of a p -group G and furthermore α_G is injective on a specific summand $\mathbf{Z}x \subset A(G)$, then $\hat{\alpha}_G$ is injective. It is the maps $\alpha_G(nx)$ that induce nontrivial maps in \mathbf{Z}/p -homology. We show even more: one gets a host of homologically distinct elements $B((\mathbf{Z}/p)^n) \rightarrow \text{cok } J_p$ for $n \geq 2$.

THEOREM D. *If $G = (\mathbb{Z}/p)^n$, and $\hat{A}_0(G) = \text{Ker}(\hat{A}(G) \rightarrow \hat{R}(G))$, then $\hat{\alpha}_G$ maps $\hat{A}_0(G)$ ($2\hat{A}_0(G)$ if $p=2$) injectively into $[BG, \text{cok } J_p]$.*

We note that Theorems A and B combine to show that $\hat{\alpha}_G$ is injective for the groups with cyclic Sylow subgroups. They are all metacyclic. Theorem C enlarges the class of groups for which $\hat{\alpha}_G$ is injective to include e.g. A_5 , the alternating group on 5 letters.

The smallest groups we cannot settle with our method are $\mathbb{Z}/4 \times \mathbb{Z}/2$, the dihedral D_8 and the quaternionic Q_8 of order 8. For these groups the maps $\alpha_G(nx)$ induce zero both in homology and K -theory. One wonders if connective K -theory or unitary bordism theory could settle these cases.

The representation ring $R(G)$ and K -theory $K(X)$ admit the structure of a λ -ring. Atiyah, Tall and Segal [3], [4] have explored the algebraic nature of such rings showing that one gets exponential isomorphisms $\varrho_k: \hat{I}(G) \xrightarrow{\cong} 1 + \hat{I}(G)$ for any p -group G , and $KSO(X)^\wedge \xrightarrow{\cong} (1 + KSO(X))^\wedge$ for any finite complex X , where $\hat{}$ denotes the p -adic completion.

Now $A(G)$ has also λ -operations, yielding λ -operations on stable cohomotopy $\pi_S^0(X) = [X, QS^0]$, see [19]. Unfortunately the λ -ring $A(G)$ is not "special", and this breaks down the algebraic program above. However, it is interesting to identify the maps $\varrho_k: Q_0S^0 \rightarrow SG_p$. We give a character argument to show

THEOREM E. $\varrho_k: Q_0S^0 \rightarrow SG_p$ is the composition $Q_0S^0 \xrightarrow{e} J_p \xrightarrow{\alpha_p} SG_p$.

The paper is divided into 4 sections. The first contains generalities on the Burnside ring: its characters, functorial properties and topology. The main theorem is 1.15 which shows that $\hat{A}(G)$ is detected by p -groups.

In section 2 we study the λ -ring structure of $A(G)$ and note that the natural map $A(G) \rightarrow R(G)$ is a λ -homomorphism. We describe the characters of the associated operations λ^n , ψ^n and ϱ_k , we show they induce operations in the (zero degree) stable cohomotopy π_S^0 (2.12) and prove Theorem E (2.20). The characters of λ^n and ψ^n were obtained independently by C. Siebeneicher [20].

The third section is devoted to the study of $\hat{\alpha}_G$ and contains the proofs of theorems A and B (3.2 and 3.3). We set up the induction machinery needed to prove theorems C and D in section 4 (4.15, 4.22 and 4.23).

The author wishes to express his gratitude to Ib Madsen, who introduced him to the subject, and helped in all possible ways. He also benefited from discussions with Jørgen Tornehave.

1. The Burnside ring.

Let G be a finite group. A G -action on a finite set S is a homomorphism from

G to Σ_S , the permutation group of S . A G -set is a finite set with a G -action. Two G -sets S and T are isomorphic if there exists a G -equivariant bijection $f: S \rightarrow T$, or in other words if the diagram

$$\begin{array}{ccc} & & \Sigma_S \\ & \nearrow & \downarrow \Sigma_f \\ G & & \Sigma_T \\ & \searrow & \end{array}$$

commutes. The equivalence classes of G -sets form a commutative semiring $A^+(G)$ under disjoint union and cartesian product. The associated ring $A(G)$ is called the Burnside ring of G .

The additive structure of $A(G)$ is easily described. Breaking G -sets into G -orbits one sees that $A(G)$ is a free abelian group with basis consisting of cosets G/H , one for each conjugacy class of subgroups H of G . We fix a set $C(G)$ of representatives.

For the multiplicative structure we introduce characters, following W. Burnside (who called them marks [5 p. 236]). Let H be a subgroup of G and S be a G -set. Then we set

$$(1.1) \quad \chi_H(S) = |S^H|,$$

the number of elements in S fixed by H . The character χ_H extends to give a ring homomorphism

$$\chi_H: A(G) \rightarrow \mathbf{Z}.$$

On the additive generators of $A(G)$ we have

$$\chi_{H_1}(G/H_2) = |\{gH_2 \mid H_1gH_2 = gH_2\}| = |\{gH_2 \mid g^{-1}H_1g \subset H_2\}|$$

or

$$(1.2) \quad \chi_{H_1}(G/H_2) = |\{gH_2 \mid H_1^g \subset H_2\}|.$$

This shows that χ_H depends only on the conjugacy class of H .

The homomorphisms χ_H define together a homomorphism

$$\chi: A(G) \rightarrow \bigoplus_{C(G)} \mathbf{Z}.$$

The first statement of the following theorem is due to W. Burnside [5] and the second one to A. Dress (unpublished, cf. [7]).

THEOREM 1.3. *The homomorphism $\chi: A(G) \rightarrow \bigoplus_{C(G)} \mathbf{Z}$ is an embedding with*

finite cokernel. Its image consists of families $(d_H)_{H \in C(G)}$ satisfying the congruences

$$d_H \equiv - \sum_{\substack{H < K \leq G \\ K/H \text{ cyclic} \neq 1}} \varphi(|K/H|)d_K \pmod{|N_G(H)/H|}$$

where we set $d_K = d_{K'}$ for $K \sim K' \in C(G)$ and φ is the Euler function.

PROOF. To prove the injectivity, it is enough to show that non-isomorphic G -sets S and T cannot have the same characters. Write $S = \sum m_H G/H$, $T = \sum n_H G/H$ with H running over $C(G)$ and let $H_0 \in C(G)$ be maximal with respect to $m_{H_0} \neq n_{H_0}$. Note from (1.2) that $\chi_{H_1}(G/H_2)$ is non-zero if and only if H_1 is conjugate to a subgroup of H_2 , denoted by $H_1 \lesssim H_2$. Then

$$\chi_{H_0}(S) = m_{H_0} \chi_{H_0}(G/H_0) + \sum_{H_0 \lesssim H, H \neq H_0} m_H \chi_{H_0}(G/H)$$

and

$$\chi_{H_0}(T) = n_{H_0} \chi_{H_0}(G/H_0) + \sum_{H_0 \lesssim H, H \neq H_0} n_H \chi_{H_0}(G/H)$$

are different as the sum terms coincide but $m_{H_0} \neq n_{H_0}$.

Let S be a G -set. We want to show that the numbers $\chi_H(S)$ satisfy the congruences for each subgroup H . If $H < K$ then the K -fixed points of S are contained in the $N_G(H)/H$ -set S^H so that we are reduced to the case $H = e$. By a theorem of W. Burnside [5 p. 191] $|G|^{-1} \sum_{g \in G} |S^g|$ is an integer, namely the number of G -orbits in S . This implies the congruence

$$|S| = \chi_e(S) \equiv - \sum_{g \neq 1} \chi_{\langle g \rangle}(S) = - \sum_{K \leq G \text{ cyclic} \neq 1} \varphi(|K|) \chi_K(S) \pmod{|G|}$$

where $\varphi(|K|)$ is the number of generators in K .

Conversely, we construct $x \in A(G)$ with given characters (d_H) by adding increasing orbits. Start with d_G points with trivial G -action. Choose a total order on $C(G)$ extending \lesssim . Assume we have $y \in A(G)$ such that $\chi_H(y) = d_H$ for H greater than H_0 . Since the characters of y and the numbers (d_H) both fulfil the congruences, we have

$$\chi_{H_0}(y) \equiv d_{H_0} \pmod{|N_G(H_0)/H_0|},$$

say $d_{H_0} = \chi_{H_0}(y) + n|N_G(H_0)/H_0|$. We add n copies of G/H_0 to y . This does not change the earlier adjusted characters, but

$$\chi_{H_0}(y + nG/H_0) = \chi_{H_0}(y) + n|N_G(H_0)/H_0| = d_{H_0}$$

by (1.2). This completes the induction.

Finally χ has finite cokernel since $\bigoplus |G|Z \subset \text{Im } \chi$ by the congruences. The theorem is proved.

REMARK. In [17] G. Segal defined a ring ω_0^0 in terms of equivariant stable homotopy. It coincides with $A(G)$ as both are characterized by theorem 1.3. For a proof and generalization to compact Lie groups, see [16, Theorem 3].

If $f: H \rightarrow G$ is a homomorphism of finite groups, then the pull-back f^*S of a G -set S has the same underlying set with H -action

$$H \xrightarrow{f} G \rightarrow \Sigma_S .$$

The induced maps $f^*: A(G) \rightarrow A(H)$ make A into a contravariant functor. The characters of f^*x are

$$(1.4) \quad \chi_U(f^*x) = \chi_{f(U)}(x), \quad U \leq H, x \in A(G) .$$

In the special case of a subgroup $i: H \rightarrow G$ we call i^* the restriction homomorphism, and denote it by Res_H^G .

There is also a covariant induction homomorphism Ind_H^G or f_* for inclusions of subgroups $f: H \rightarrow G$. On the coset basis it is given by

$$\text{Ind}_H^G(H/H_1) = G/H_1, \quad H_1 \leq H \leq G$$

It is easily checked from (1.2) that the characters of f_*y are

$$(1.5) \quad \chi_U(f_*y) = \sum_{U^* \leq H} \chi_{U^*}(y), \quad U \leq G, y \in A(H)$$

where g runs through representatives of G/H .

The homomorphisms Res and Ind are related in the same fashion as the restriction and induction maps in representation theory or cohomology of finite groups. If H is a subgroup of G , then the Frobenius reciprocity

$$(1.6) \quad \text{Ind}_H^G(y \cdot \text{Res}_H^G(x)) = \text{Ind}_H^G(y) \cdot x, \quad y \in A(H), x \in A(G)$$

holds. Further, if K is another subgroup of G , let $Kg_1H, \dots, Kg_rH \subset G$ be the double cosets of $G \bmod (K, H)$. Then we have

$$(1.7) \quad \text{Res}_K^G \text{Ind}_H^G(x) = \sum_{i=1}^r \text{Ind}_{K \cap g_i H g_i^{-1}}^K (c_{g_i}^* \text{Res}_{g_i^{-1} K g_i \cap H}^H(x)), \quad x \in A(H)$$

where c_{g_i} is conjugation by g_i . The proofs of (1.6) and (1.7) are analogous to the corresponding formulas of representation theory.

Each G -set S can be considered as a linear representation of G over a field k by extending the G -action on the canonical basis of k^S by linearity. As the trace of a permutation matrix is equal to the number of 1's along the diagonal, the

linear character of k^S can be read off from the characters χ_H of (1.1) for H cyclic:

$$\chi_{k^S}(g) = \chi_{\langle g \rangle}(S).$$

If k has characteristic 0, then the elements in $R_k(G)$ are detected by their linear characters. We conclude from theorem 1.3

LEMMA 1.8. *Let k be a field of characteristic 0. Then the kernel of the natural map $A(G) \rightarrow R_k(G)$ coincides with the kernel of the restriction map*

$$\text{Res}: A(G) \rightarrow \bigoplus_{C \leq G} A(C)$$

to the cyclic subgroups of G .

In Segal's conjecture the Burnside ring $A(G)$ is compared with the stable cohomotopy ring $\pi_S^0(BG)$. As the latter is complete (see section 3), we study here the algebraic process of completing $A(G)$.

The character χ_e just counts the number of points in a G -set and defines an augmentation $\varepsilon: A(G) \rightarrow \mathbb{Z}$. This is a split surjection so $A(G) = \mathbb{Z} \oplus I(G)$ where $I(G)$ is the augmentation ideal $\varepsilon^{-1}(0)$.

We give the ring $A(G)$ the usual $I(G)$ -adic topology, letting the powers $I(G)^n$ be a neighbourhood basis of 0. The completion of $A(G)$ is defined as the inverse limit

$$\hat{A}(G) = \lim_{\leftarrow n} A(G)/I(G)^n.$$

We shall study the kernel of $A(G) \rightarrow \hat{A}(G)$. First we recall a result from commutative algebra (see e.g. [22 p. 262, Corollary to Theorem 8]). Let A be a Noetherian ring with no nilpotents and $\mathfrak{m} \subset A$ a prime ideal. Then the kernel of the natural map from A to the \mathfrak{m} -adic completion $\hat{A} = \lim_{\leftarrow} A/\mathfrak{m}^n$ is

$$\bigcap_{n=0}^{\infty} \mathfrak{m}^n = \bigcap_{\mathfrak{p}_j + \mathfrak{m} \neq A} \mathfrak{p}_j$$

where \mathfrak{p}_j runs over such minimal prime ideals of A that $\mathfrak{p}_j + \mathfrak{m} \neq A$.

The ring $A(G)$ is Noetherian as a finitely generated abelian group. A. Dress determined in [7] the prime ideal structure of $A(G)$. There are two types of prime ideals in $A(G)$: the minimal ones

$$\mathfrak{p}_{U,0} = \{x \in A(G) \mid \chi_U(x) = 0\}$$

for $U \leq G$, and the maximal ones

$$\mathfrak{p}_{U,p} = \{x \in A(G) \mid \chi_U(x) \equiv 0 \pmod{p}\}$$

for $U \leq G$ and p a prime. Furthermore,

$$(1.9) \quad \begin{aligned} \mathfrak{p}_{U,0} &= \mathfrak{p}_{V,0} && \text{if and only if } U \sim V, \\ \mathfrak{p}_{U,p} &= \mathfrak{p}_{V,q} && \text{if and only if } p=q \text{ and } U^p \sim V^p \end{aligned}$$

where U^p is the smallest normal subgroup of U with U/U^p a p -group, and $\mathfrak{p}_{U,0} \subset \mathfrak{p}_{U,p}$ together with (1.9) accounts for all inclusions between prime ideals in $A(G)$.

PROPOSITION 1.10. *The kernel of $A(G) \rightarrow \hat{A}(G)$ coincides with the kernel of the restriction map*

$$\text{Res: } A(G) \rightarrow \bigoplus_{G_p \leq G} A(G_p)$$

to the Sylow subgroups G_p of G .

PROOF. It follows from the above that the kernel is

$$\bigcap_{n=0}^{\infty} I(G)^n = \bigcap_{\mathfrak{p}_{U,0} + I(G) \neq A(G)} \mathfrak{p}_{U,0}.$$

Now $I(G) = \mathfrak{p}_{e,0}$ and if the ideal $\mathfrak{p}_{U,0} + \mathfrak{p}_{e,0}$ is proper then it is contained in a maximal ideal $\mathfrak{p}_{V,p}$. By (1.9) this implies that $\mathfrak{p}_{U,p} = \mathfrak{p}_{e,p} = \mathfrak{p}_{V,p}$ hence $U^p = e = V^p$ and U is a p -group. Conversely, if U is a p -group then $\mathfrak{p}_{U,0} + \mathfrak{p}_{e,0} \subset \mathfrak{p}_{U,p}$. Thus

$$\text{Ker } (A(G) \rightarrow \hat{A}(G)) = \bigcap_{U \leq G \text{ } p\text{-group}} \text{Ker } \chi_U,$$

and the claim follows from theorem 1.3.

COROLLARY 1.11. *If G is a p -group then $A(G) \rightarrow \hat{A}(G)$ is a monomorphism.*

In the case of a p -group G the $I(G)$ -adic completion is the familiar p -adic one:

$$\hat{A}(G) = \mathbf{Z} \oplus (\hat{\mathbf{Z}}_p \otimes_{\mathbf{Z}} I(G))$$

where $\hat{\mathbf{Z}}_p = \varprojlim_n \mathbf{Z}/p^n$ denotes the p -adic integers:

PROPOSITION 1.12. *If G is a p -group then the $I(G)$ -adic topology of $A(G)$ is the same as its p -adic topology.*

PROOF. We have to prove that for each m there are integers n_1, n_2 such that

$$(1) \quad p^{n_1} I(G) \subset I(G)^m$$

$$(2) \quad I(G)^{n_2} \subset p^m I(G).$$

The first relation follows from Atiyah's

LEMMA 1.13. *For any group G , $|G|I(G)^n \subset I(G)^{n+1}$.*

(This is a consequence of the reciprocity formula 1.6, see [1p. 269, Proposition 6.13]).

To get the inclusion (2), we note that for any $H \leq G$ and $U \leq G$

$$\chi_U(G/H - |G/H|) \equiv 0 \pmod{p}$$

since the complement of $(G/H)^U$ consists of non-trivial U -orbits, hence $\chi_U(I(G)) \subset p\mathbb{Z}$. As $\chi = \bigoplus_{U \leq G} \chi_U$ is a ring homomorphism we have $\chi(I(G)^n) \subset \bigoplus_{e \neq U \leq G} p^n \mathbb{Z}$ and it is enough to prove

$$\bigoplus_{e \neq U \leq G} |G| \mathbb{Z} \subset \chi(I(G)).$$

This follows immediately from the congruences 1.3. The proof of 1.12 is complete.

The completion $\hat{A}(G_p)$ is now described for p -groups G_p . Next we shall show that if G is an arbitrary finite group, then $\hat{A}(G)$ embeds into the sum $\bigoplus \hat{A}(G_p)$, taken over the Sylow subgroups G_p of G . This is done by completing the map of proposition 1.10.

Let A be a Noetherian ring and $\mathfrak{m} \subset A$ an ideal. The \mathfrak{m} -adic completion of a finitely generated A -module M is defined to be $\hat{M} = \varprojlim_n M/\mathfrak{m}^n M$. It is a basic fact that Noetherian completion is an exact functor [1 p. 258, Proposition 3.16].

If $H \leq G$, then $A(H)$ is an $A(G)$ -module via the restriction homomorphism $\varrho = \text{Res}_H^G: A(G) \rightarrow A(H)$. In the following proof we distinguish the prime ideals $\mathfrak{p}_{U,p}$ of $A(H)$ and $A(G)$ by upper indices, so that $\mathfrak{p}_{U,p}^H \subset A(H)$ and $\mathfrak{p}_{U,p}^G \subset A(G)$.

PROPOSITION 1.14. *Let H be a subgroup of G . Then the $I(H)$ -adic topology of $A(H)$ is the same as its $I(G)$ -adic topology.*

PROOF. It is enough to show that the radicals of the ideals $\varrho(I(G))$ and $I(H)$ coincide [22 p. 256]. This means that each prime ideal $\mathfrak{p} \subset A(H)$ either contains the both ideals or none. Since $\varrho(I(G)) \subset I(H)$, one way is trivial. Let \mathfrak{p} be a prime ideal of $A(H)$ with $\varrho(I(G)) \subset \mathfrak{p}$. We claim that $I(H) \subset \mathfrak{p}$. We know that \mathfrak{p} is of the form $\mathfrak{p}_{U,0}^H$ or $\mathfrak{p}_{U,p}^H$ with some subgroup $U \leq H$ and some prime p . If $\mathfrak{p} = \mathfrak{p}_{U,0}^H$, then

$$\mathfrak{p}_{e,0}^G = I(G) \subset \varrho^{-1}(\mathfrak{p}) = \mathfrak{p}_{U,0}^G$$

which implies $U = e$ and $\mathfrak{p} = \mathfrak{p}_{e,0}^H = I(H)$ by (1.9). Similarly, if $\mathfrak{p} = \mathfrak{p}_{U,p}^H$, then

$$\mathfrak{p}_{e,0}^G = I(G) \subset \varrho^{-1}(\mathfrak{p}) = \mathfrak{p}_{U,p}^G$$

and U must be a p -group by (1.9), whence $\mathfrak{p} = \mathfrak{p}_{U,p}^H = \mathfrak{p}_{e,p}^H$. In both cases $I(H) \subset \mathfrak{p}$, are claimed.

THEOREM 1.15. *Let G be a finite group and $\{G_p\}$ its Sylow subgroups. Then the completion of the restriction maps $\text{Res}_{G_p}^G$ gives an injective homomorphism*

$$0 \rightarrow \hat{A}(G) \rightarrow \bigoplus_p \hat{A}(G_p).$$

PROOF. By 1.10 $A(G)/\bigcap_{n=0}^{\infty} I^n(G)$ maps injectively into $\bigoplus_p A(G_p)$. Both modules have $I(G)$ -adic topology by 1.14. The claim follows since Noetherian completion is an exact functor.

We close the chapter with some examples. The first two are abelian p -groups. The last one illustrates the restrictions to Sylow subgroups and completion.

EXAMPLE 1.16. The cyclic group \mathbb{Z}/p^n . It has a unique subgroup of order p^{n-m} , $0 \leq m \leq n$. Let η_m be the quotient ($\eta_0 = 1$). We have additively

$$A(\mathbb{Z}/p^n) = \mathbb{Z} \oplus \mathbb{Z}\eta_1 \oplus \dots \oplus \mathbb{Z}\eta_n.$$

From the characters

$$\chi_{\mathbb{Z}/p^i}(\eta_m) = \begin{cases} p^m, & i \leq n-m \\ 0, & i > n-m \end{cases}$$

one gets the multiplication $\eta_1 \cdot \eta_m = p^l \eta_m$ for $l \leq m$.

EXAMPLE 1.17. The elementary abelian group $(\mathbb{Z}/p)^n$. It can be interpreted geometrically as a vector space over the finite field F_p , with subgroups corresponding to linear subspaces. The number of m -dimensional planes is


$$\begin{aligned} G(m, n) &= \frac{(p^n - 1)(p^n - p) \dots (p^n - p^{m-1})}{(p^m - 1)(p^m - p) \dots (p^m - p^{m-1})} \\ &= \frac{(p^n - 1)(p^{n-1} - 1) \dots (p^{n-m+1} - 1)}{(p^m - 1)(p^{m-1} - 1) \dots (p - 1)} \end{aligned}$$

(G stands for Grassmann). $A((\mathbb{Z}/p)^n)$ is additively generated by the m -

dimensional quotient planes η_m^i , $0 \leq m \leq n$, $1 \leq i \leq G(n-m, n) = G(m, n)$, and

$$F_p^n/V_1 \times F_p^n/V_2 = |F_p^n/V_1 + V_2|F_p^n/V_1 \cap V_2.$$

EXAMPLE 1.18. The alternating group A_4 . The diagram of subgroups is



and the character table is given in table 1.19 where the small letters 1, a, b, c, and d denote the cosets A_4/H in the given order.

Table 1.19.

$A_4/H \backslash \chi_H$	A_4	$Z/2 \oplus Z/2$	$Z/3$	$Z/2$	e
1	1	1	1	1	1
a	0	3	0	3	3
b	0	0	1	0	4
c	0	0	0	2	6
d	0	0	0	0	12

The Burnside rings of the Sylow subgroups of A_4 are described in the preceding examples:

$$A(Z/2 \oplus Z/2) = Z \oplus Z\eta_1^1 \oplus Z\eta_1^2 \oplus Z\eta_1^3 \oplus Z\eta_2$$

$$A(Z/3) = Z \oplus Z\xi.$$

The restriction map $A(G) \rightarrow A(G_2) \oplus A(G_3)$ is read from the character table using (1.4) $\chi_H(i^*x) = \chi_H(x)$. The result is

$$\begin{array}{ll} 1 \rightarrow (1, 1) & \\ a \rightarrow (3, \xi) & \\ b \rightarrow (\eta_2, 1 + \xi), & \text{or } b_1 \rightarrow (\eta_2, 0) \\ c \rightarrow (\eta_1^1 + \eta_1^2 + \eta_1^3, 2\xi) & c_1 \rightarrow (\eta_1^1 + \eta_1^2 + \eta_1^3, 0) \\ d \rightarrow (3\eta_2, 4\xi) & d_1 \rightarrow (0, 0) \end{array}$$

in the basis $a_1 = a - 3$, $b_1 = b - a - 1$, $c_1 = c - 2a$, $d_1 = d - 3b - a + 3$ for $I(A_4)$. Here \bar{x} denotes $x - \varepsilon(x) \in I(G_p)$. This shows that the image of $A(G)$ in $A(G_2) \oplus A(G_3)$ consists precisely of the *stable* elements. These are the pairs (x_2, x_3) with

- (1.20) (1) $\varepsilon(x_2) = \varepsilon(x_3)$
 (2) $\chi_{H_1}(x_p) = \chi_{H_2}(x_p)$, if $H_1 < G_p$ and $H_2 < G_p$ are conjugate in G .

The condition (2) rules out η_1^1, η_1^2 and η_1^3 since they have different characters on the three $\mathbb{Z}/2 < \mathbb{Z}/2 \oplus \mathbb{Z}/2$ which are conjugate in A_4 .

It might be interesting to know whether (1.20) characterizes the image of $A(G)$ in $\bigoplus_p A(G_p)$ in general. If the Segal conjecture $\hat{A}(G) = \pi_0^S(BG)$ is true, then this holds at least on the completion level by general properties of cohomology theories on BG (see [11, 1.7]).

Finally, the multiplication table for $I(A_4)$

	a_1	b_1	c_1	d_1
a_1	$-3a_1$	d_1	0	$-3d_1$
b_1		$-4b_1 - d_1$	$-4c_1$	$-d_1$
c_1			$6b_1 - 10c_1 + 2d_1$	0
d_1				$3d_1$

shows that $\hat{A}(G) = \mathbb{Z} \oplus \hat{\mathbb{Z}}_3 a_1 \oplus \hat{\mathbb{Z}}_2 b_1 \oplus \hat{\mathbb{Z}}_2 c_1$ and $I(G)^\infty = \mathbb{Z} d_1$.

2. λ -Operations on the Burnside ring.

Let G be a finite group. If k is a field and V is a representation of G over k then the exterior powers $\lambda^n V$ are also G -representations. We want to construct operations in $A(G)$ which under the natural map $A(G) \rightarrow R_k(G)$ correspond to the exterior powers. As it is not clear how to make sense of the relation $x \wedge y = -y \wedge x$ in a G -set, we consider first the symmetric powers $s^n V$ where no signs are needed.

Let $S = \{s_1, \dots, s_l\}$ be a G -set. The vector space $s^n(k^S)$ has a basis consisting of monomials of degree n in $s_i \in s^1(k^S)$, considered as elements of the symmetric algebra $s(k^S)$. We define the n th symmetric power of S as

$$s^n(S) = S^n / \Sigma_n$$

with the diagonal G -action. It is clear that

(2.1)
$$s^1(S) = S$$

(2.2)
$$s^n(S \cup T) = \sum_{i=0}^n s^i(S) s^{n-i}(T).$$

We assign to S the formal power series

$$(2.3) \quad s_t(S) = 1 + \sum_{n \geq 1} s^n(S)t^n \in A(G)[[t]] .$$

It is invertible as the leading coefficient is 1, and (2.2) shows that $s_t(S \cup T) = s_t(S) \cdot s_t(T)$. The homomorphism s_t is uniquely extended to $A(G)$ by $s_t(S - T) = s_t(S) \cdot s_t(T)^{-1}$.

In the representation ring $R_k(G)$ the symmetric powers are connected to the exterior powers by means of the identity

$$\lambda_t(V)s_{-t}(V) = 1 .$$

Thus we are lead to

DEFINITION 2.4. The n th exterior power of $x \in A(G)$, denoted by $\lambda^n(x)$, is the coefficient of t^n in the series $\lambda_t(x) = s_{-t}(x)^{-1}$.

The formulae (2.1)–(2.3) translate to give

$$(2.5) \quad \begin{aligned} \text{(i)} \quad & \lambda^0(x) = 1 \\ \text{(ii)} \quad & \lambda^1(x) = x \\ \text{(iii)} \quad & \lambda^n(x + y) = \sum_{i=0}^n \lambda^i(x)\lambda^{n-i}(y) . \end{aligned}$$

This is summarised in saying that the operations $\lambda^n, n \geq 1$, give $A(G)$ the structure of a λ -ring [4]. By construction they are natural with respect to induced maps, and $A(G) \rightarrow R_k(G)$ is a λ -homomorphism.

We shall calculate the character of $\lambda^n(x), \chi_U(\lambda^n(x))$. By (2.5) and naturality it is enough to consider a single G -orbit $x = G/H$. A point $(s_1, \dots, s_n) \in (G/H)^n/\Sigma_n$ is fixed under G only if it can be split up to G -orbits G/H . This implies that n is a multiple of $|G/H|$, and

$$(2.6) \quad \chi_G(s^n(G/H)) = \begin{cases} 0, & n \not\equiv 0 \pmod{|G/H|} \\ 1, & n \equiv 0 \pmod{|G/H|} . \end{cases}$$

Thus $\chi_G(s_t(G/H)) = (1 - t^{|G/H|})^{-1}$. For a subgroup U of G we can break up S into U -orbits $\cup S_i$ so

$$(2.7) \quad \chi_U(\lambda_t(S)) = \prod_{S_i \subset S} \prod_{U\text{-orbits}} (1 - (-t)^{|S_i|}) .$$

In particular the degree of $\chi_U(\lambda_t(S))$ is equal to $|S|$, hence $\lambda^n(S) = 0$ if $n > |S|$. Also $\varepsilon(\lambda_t(S)) = \chi_e(\lambda_t(S)) = (1 + t)^{|S|}$. Thus $A(G)$ is a finite-dimensional augmented λ -ring.

We define the Adams operations $\psi^n: A(G) \rightarrow A(G)$, $n \geq 1$, by

$$\psi_{-t}(x) = -t \frac{\lambda'_t(x)}{\lambda_t(x)}, \quad \text{where } \psi_t(x) = \sum_{n \geq 1} \psi^n(x)t^n.$$

Then (2.5) implies that ψ^n is additive, $\psi^n(x+y) = \psi^n(x) + \psi^n(y)$. As to the characters, the logarithmic differentiation of (2.7) yields

$$\chi_U(\psi_t(X)) = \sum_{S_i \in S} \sum_{U\text{-orbits}} |S_i| t^{|S_i|} (1 - t^{|S_i|})^{-1}.$$

This proves

PROPOSITION 2.8. $\chi_U(\psi^n(S)) = \sum_{|S_i| \mid n} |S_i|$, where $S = \bigcup S_i$ the decomposition into U -orbits.

COROLLARY 2.9. *The Adams operations are periodic of period dividing the order of G .*

(Indeed, the length of each U -orbit U/H is a divisor of $|G|$.)

If G is a p -group and $(n, p) = 1$, then the only orbits occurring in 2.8 are the U fixed points. This proves

COROLLARY 2.10. *If G is a p -group, then $\psi^n = \text{id}$ for n relatively prime to p .*

The operations λ^n have geometrical significance: they induce natural transformations of π_S^0 , the zeroth stable cohomotopy functor.

First recall the Barratt–Quillen theorem. The group completion map $i: \coprod_{n \geq 1} B\Sigma_n \rightarrow QS^0$ gives a natural transformation of monoid-valued functors

$$\left[X, \coprod_{n \geq 1} B\Sigma_n \right] \rightarrow [X, QS^0].$$

Here $A(X) = [X, \coprod_{n \geq 1} B\Sigma_n]$ is the set of isomorphism classes of finite coverings of X organized to a semiring under disjoint union and fibrewise cartesian product of the total spaces, and $\pi_S^0(X) = [X, QS^0]$ is the stable cohomotopy of X (in degree 0), an abelian group with respect to loop sum. The group completion theorem states [18, Proposition 4.1]

THEOREM 2.11. *The transformation $A \rightarrow \pi_S^0$ is universal among transformations $\theta: A \rightarrow F$, where F is a representable abelian-group-valued homotopy functor on compact spaces, and θ is a transformation of monoid-valued functors.*

We deduce from this the existence of λ -operations on π_S^0 along the lines of Segal [19].

THEOREM 2.12. *There are natural transformation $\lambda^n: \pi_S^0 \rightarrow \pi_S^0$ for $n \geq 0$, such that*

- (i) $\lambda^0(x) = 1$
- (ii) $\lambda^1(x) = x$
- (iii) $\lambda^n(x + y) = \sum_{i=0}^n \lambda^i(x) \lambda^{n-i}(y)$.

PROOF. We define a transformation $\lambda^n: A(X) \rightarrow \pi_S^0(X)$. Assume X is connected. An m -fold covering $Y \downarrow X$ can be written as $P \times_{\Sigma_m} [m] \downarrow X$, where P is the principal Σ_m -bundle associated to Y consisting of mappings of $[m] = (1, 2, \dots, m)$ onto the fibres of Y , and $[m]$ has the usual Σ_m -action. Let $\lambda^n([m]) = S - T \in A(\Sigma_n)$. We associate to $Y \downarrow X$ the difference

$$\lambda^n(Y) = P \times_{\Sigma_m} S - P \times_{\Sigma_m} T \in \pi_S^0(X)$$

where we have used $A \rightarrow \pi_S^0$ from 2.11.

Let us form the mapping

$$\lambda_t = \sum_{n \geq 0} \lambda^n t^n: A(X) \rightarrow 1 + \pi_S^0(X)[[t]]^+ = \prod_{n \geq 1} \pi_S^0(X).$$

It is a monoid homomorphism, when we use multiplication of power series on the right. As $1 + \pi_S^0(X)[[t]]^+$ is a representable abelian-group-valued functor, λ_t extends by theorem 2.11 to a group homomorphism

$$\lambda_t: \pi_S^0(X) \rightarrow 1 + \pi_S^0(X)[[t]]^+.$$

This completes the proof of theorem 2.12.

In the articles [3] and [4] Atiyah, Tall and Segal showed that special p -adic λ -rings possess certain canonical exponential isomorphisms between the additive group $\hat{I}(G)$ and the multiplicative group $1 + \hat{I}(G)$. Unfortunately the Burnside ring $A(G)$ is special only if G is cyclic: The Adams operations ψ^n are ring homomorphisms in special λ -rings, but Siebeneicher showed that this is not true in $A(G)$ for any non-cyclic G [20, p. 232]. On the other hand, $A(G)$ embeds as a sub- λ -ring of $R(G)$ if G is cyclic.

However, it is interesting to study the exponential map ϱ_k . We first do the algebra and then identify the resulting geometric map $Q_0 S^0 \rightarrow SG[1/k]$ with the composition

$$Q_0 S^0 \xrightarrow{e} \text{Im } J_p \xrightarrow{a_p} SG \left[\frac{1}{k} \right].$$

Philosophically this is a negative result: the λ -operations on $A(G)$ do not give any information on the fibre of e , the space usually denoted $\text{cok } J_p$.

Let G be a finite group. We shall encounter series of the form $\lambda_\alpha(x) = 1 + \sum \alpha^n \lambda^n(x)$. To show their convergence in $\hat{A}(G)$ we introduce a new topology on $A(G)$. Define the Grothendieck operations by

$$(2.13) \quad \gamma^n(x) = \lambda^n(x + n - 1).$$

If $\gamma_t(x) = 1 + \sum_{n \geq 1} \gamma^n(x) t^n$, then

$$(2.14) \quad \gamma_t(x) = \lambda_{t/(1-t)}(x), \quad \gamma_t(x+y) = \gamma_t(x)\gamma_t(y).$$

The γ -operations are convenient on the augmentation ideal $I(G)$ as the generators $G/H - \varepsilon(G/H)$ have finite γ -dimension but infinite λ -dimension. In fact, (2.7) and (2.14) imply

$$\chi_U(\gamma_t(S - \varepsilon(S))) = \prod_{S_i \in S \text{ U-orbit}} [(1-t)^{|S_i|} - (-t)^{|S_i|}]$$

for any G -set S .

Define the γ -filtration by

$$(2.15) \quad I_n \text{ is the group generated additively by } \gamma^{n_1}(x_1) \dots \gamma^{n_r}(x_r) \\ \text{with } x_i \in I(G), \sum n_i \geq n.$$

Then $I_m \cdot I_n \subset I_{m+n}$, $I_0 = A(G)$ and $I_1 = I(G)$. Thus the filtration $(I_n)_{n \geq 0}$ defines a topology on $A(G)$, the γ -topology.

PROPOSITION 2.16. *If G is a p -group, then the p -adic, $I(G)$ -adic and γ -topologies on $A(G)$ are equivalent.*

PROOF. We proved in 1.12 that the first two topologies coincide. Atiyah [1, Corollary 12.3] shows that the γ -topology is equivalent to the $I(G)$ -adic if $I(G)$ has a finite number of generators, each of finite γ -dimension.

Let G be a p -group $x \in I(G)$ and $\alpha \in \hat{\mathbf{Z}}_p$. Then the series $\gamma_\alpha(x) = 1 + \sum_{n \geq 1} \alpha^n \gamma^n(x)$ converges in the γ -topology, hence also in $\hat{A}(G)$. More generally, if $\alpha \in A$ where A is a finitely generated $\hat{\mathbf{Z}}_p$ -algebra, then $\gamma_\alpha(x)$ exists in $1 + \hat{I}(G) \otimes_{\hat{\mathbf{Z}}_p} A$. We fix a prime k different from p and apply this to $A = \hat{\mathbf{Z}}_p[\xi]$, where ξ is a primitive k th root of 1.

DEFINITION 2.17. $\varrho_k(x) = \prod_{u \neq 1} \lambda_{-u}(x)$, $x \in I(G)$.

A priori $\varrho_k(x)$ belongs to $1 + \hat{I}(G) \otimes \hat{\mathbf{Z}}_p[\xi]$. But it is invariant under the action of the Galois group of $\hat{\mathbf{Q}}_p(\xi)/\hat{\mathbf{Q}}_p$, so actually $\varrho_k(x) \in 1 + \hat{I}(G)$.

We compute the character of $\varrho_k(x)$. A substitution in (2.7) yields

$$\chi_U(\varrho_k(S - \varepsilon(S))) = \prod_{\substack{u^k=1 \\ u \neq 1}} \left(\prod_{S_i \subset S} (1 - u^{|S_i|}) \right) (1 - u)^{-|S|}.$$

As the sizes of the U -orbits S_i are 1 or multiples of p and $(k, p) = 1$, $1 - u^{|S_i|}$ runs through the same values as $1 - u$, when S_i is fixed. Noting that $\prod_{\substack{u^k=1 \\ u \neq 1}} (1 - u) = k$ we get

$$(2.18) \quad \chi_U(\varrho_k(S - \varepsilon(S))) = k^{o_U(S) - \varepsilon(S)}$$

where $o_U(S)$ is the number of U -orbits in S .

Next we show that ϱ_k can be obtained by a direct operation on G -sets.

PROPOSITION 2.19. *For a G -set S let $\theta_k(S)$ be the underlying set of the vector space F_k^S with the linear G -action extending the permutation of the basis. Then θ_k satisfies*

(i) $\theta_k(S + T) = \theta_k(S)\theta_k(T)$

(ii) $\varepsilon\theta_k(S) = k^{\varepsilon(S)}$

(iii) θ_k is natural

(iv) $\theta_k(S) = \prod_{\substack{u^k=1 \\ u \neq 1}} \lambda_{-u}(S)$

on $A^+(G)$.

PROOF. Properties (i)–(iii) are obvious. To prove (iv) it is enough to check $\chi_U(\theta_k(S))$ for $U = G$ by naturality and for a transitive G -set S by (i). A point $\sum_{x \in S} a_x x$, $a_x \in F_k$, is fixed under G only if it is of the form $a \sum_{x \in S} x$, thus

$$\chi_G(\theta_k(S)) = |F_k| = k$$

but $\chi_G(\prod \lambda_{-u}(S)) = k^{o_G S} = k$.

REMARK. There is no problem about the convergence of $\lambda_{-u}(S)$ in (iv), since $\lambda_t(S)$ is a polynomial.

We return now to the stable cohomotopy interpretation. Let p and k be different primes. The operation $\theta_k: A^+(\Sigma_n) \rightarrow A^+(\Sigma_n)$ induces a natural transformation $\theta_k: A \rightarrow A$ as in theorem 2.12: if $Y \downarrow X$ is an n -fold covering, write it as $Y = P \times_{\Sigma_n} [n]$ with some principal Σ_n -bundle P and set $\theta_k(Y) = P \times_{\Sigma_n} \theta_k[n]$. By (2.19) (i) and (ii) the composite

$$A \xrightarrow{\theta_k} A \rightarrow \pi_S^0$$

is exponential and maps n -fold coverings of X to the component $[X, Q_k S^0]$.

In order to apply theorem 2.11 the elements $\theta_k(Y) \in \pi_S^0(X)$ have to be invertible in the composition product, in particular the maps in $Q_k S^0$ must have an inverse of degree k^{-n} . This can be accomplished by forming the localization $\lambda: Q S^0 \rightarrow Q S_p^0$ of the space $Q S^0$ at p [21, sections 2 and 4]. Denote the 1-component of $Q S_p^0$ by SG_p . Then the transformation

$$\varrho_k: A(X) \rightarrow [X, SG_p]$$

which takes an n -fold covering $Y \downarrow X$ to

$$X \xrightarrow{\theta_k(Y)} Q_k S^0 \xrightarrow{\lambda} Q_k S_p^0 \xrightarrow{\cdot k^{-n}} SG$$

extends to a homomorphism

$$(*) \quad \varrho_k: \pi_S^0(X) \rightarrow [X, SG_p]$$

by theorem 2.11.

The restriction of $(*)$ to $\tilde{\pi}_S^0$ corresponds to an H -map $\varrho_k: Q_0 S^0 \rightarrow SG_p$, defined up to homotopy. The space SG_p splits as a product $J_p \times \text{cok } J_p$ (this will be discussed in section 4), and we point out here

THEOREM 2.20. *The map $\varrho_k: Q_0 S^0 \rightarrow SG_p$ factors $Q_0 S^0 \xrightarrow{e} J_p \xrightarrow{\alpha_p} SG_p$.*

PROOF. Compare proposition 2.19 to [14, p. 236].

The natural homomorphism $A(G) \rightarrow R(G)$ is a λ -ring homomorphism. For the elementary abelian groups its kernel is large. We evaluate the Adams operations on $A((\mathbb{Z}/p)^n)$ in the concluding example.

EXAMPLE 2.21. Elementary abelian groups $(\mathbb{Z}/p)^n$.

Let $G = (\mathbb{Z}/p)^n$. Each generator G/H of $A(G)$ is the image of the regular representation under $\pi^*: A(G/H) \rightarrow A(G)$. By naturality it is thus enough to find $\psi^k(\eta_n)$, where we denote $\eta_n = G/e$ (see 1.17). From 2.8 we get

$$\chi_H(\psi^k(\eta_n)) = \begin{cases} 0, & k \not\equiv 0 \pmod{|H|} \\ p^n, & k \equiv 0 \pmod{|H|} \end{cases} \quad H \leq (\mathbb{Z}/p)^n$$

which depends only on the size of H . This suggests that we begin with the sum of all cosets of cardinality p^m ,

$$\eta_m^{\text{tot}} = \sum_i \eta_m^i = \sum_{|H|=p^{n-m}} G/H$$

which has the characters $\chi_H(\eta_m^{\text{tot}}) = 0$ if $|H| > p^{n-m}$, $\chi_{\mathbb{Z}/p^{n-m}}(\eta_m^{\text{tot}}) = p^m$, and then correct χ_H for smaller H by adding linear combinations of η_{m+k}^{tot} , $k > 0$. An inductive calculation shows that the element

$$(2.22) \quad a_m = \sum_{k=0}^{n-m} (-1)^k p^{k(k-1)/2} G(k, m+k) \frac{\eta_{m+k}^{\text{tot}}}{p^{m+k}} \in A(G) \left[\frac{1}{p} \right],$$

where $G(k, m+k)$ is defined in example 1.17, has characters

$$\chi_{(\mathbb{Z}/p)^k}(a_m) = \begin{cases} 1, & k = n - m \\ 0, & k \neq n - m. \end{cases}$$

By the first formula we then get

$$\psi^{p^m h}(\eta_n) = \sum_{i=n-m}^n p^i a_i = p^m \sum_{k=0}^m (-1)^k p^{\binom{k}{2}} G(k, n-m+k-1) \eta_{n-m+k}^{\text{tot}}$$

if $(p, h) = 1, 0 \leq m < n$ and

$$\psi^{p^m h}(\eta_n) = p^n.$$

3. The map α_G .

In this section we study the injectivity of the map $\hat{\alpha}_G: \hat{A}(G) \rightarrow \pi_S^0(BG)$ from the completion of the Burnside ring of a finite group G to the stable cohomology of its classifying space BG .

Recall the definition of α_G . Each G -set S with G -action $\varrho: G \rightarrow \Sigma_{|S|}$ gives rise to a map $\alpha_G(S): BG \rightarrow QS^0$ by

$$BG \xrightarrow{B\varrho_*} B\Sigma_{|S|} \hookrightarrow \coprod_{n \geq 0} B\Sigma_n \xrightarrow{i} QS^0,$$

where i is the group completion map. The homotopy class of $\alpha_G(S)$ depends only on the class of S in $A(G)$, and the correspondence $S \mapsto \alpha_G(S)$ extends to a ring homomorphism

$$\alpha_G: A(G) \rightarrow [BG, QS^0]$$

(by definition, $\pi_S^0(BG) = [BG, QS^0]$).

Alternatively, we can define $\alpha_G(S)$ as the image of the covering $EG \times_G S \downarrow BG$ in $\pi_S^0(BG)$ (cf. 2.11). This is quite analogous to the homomorphism

$$\alpha: R(G) \rightarrow K^*(BG)$$

studied by Atiyah in [1]: if $\varrho: G \rightarrow \text{Gl}(n, \mathbb{C})$ is a complex representation of G , then $\alpha(\varrho)$ is the class of the vector bundle $EG \times_G (\mathbb{C}^n, \varrho)$ in $K^0(BG)$. It is no surprise that α_G and α are connected via the natural map $A(G) \rightarrow R(G)$:

PROPOSITION 3.1. *Let G be a finite group. Then the diagram*

$$\begin{array}{ccc} A(G) & \xrightarrow{\alpha_G} & \pi_S^0(BG) \\ \downarrow & & \downarrow e_* \\ R(G) & \xrightarrow{\alpha} & K^*(BG) \end{array}$$

commutes, where $e: QS^0 \rightarrow BU \times Z$ is induced from a unit of the unitary spectrum.

PROOF. Let S be a G -set of cardinality n , and let $\varrho: G \rightarrow \Sigma_n$ be its G -action. If $P: \Sigma_n \rightarrow U_n$ is the permutation representation, then $\alpha_G(S)$ and $\alpha(\mathbb{C}^S)$ are represented by the upper and lower horizontal arrows in

$$\begin{array}{ccccccc}
 BG & \xrightarrow{B\varrho} & B\Sigma_n & \xrightarrow{i_n} & Q_n S^0 & \hookrightarrow & QS^0 \\
 \parallel & & \downarrow BP & & \downarrow e_n & & \downarrow e \\
 BG & \xrightarrow{BP \circ B\varrho} & BU_n & \longrightarrow & BU \times (n) & \hookrightarrow & BU \times Z
 \end{array}$$

But the right hand squares commute [12, Corollary 5.31].

REMARK. We shall give in section 4 a closer description of the map e (see 4.18).

In section 1 we considered the $I(G)$ -adic topology on $A(G)$. If X is a CW-complex with n -skeleton X^n , we filter $\pi_S^0(X)$ by

$$(F) \quad F^n \pi_S^0(X) = \text{Ker} (\pi_S^0(X) \rightarrow \pi_S^0(X^{n-1})).$$

Then $F^n \cdot F^m \subset F^{n+m}$ by diagonal approximation. J. W. Milnor's original construction of BG gave a CW-complex with finite skeletons $B_n G$. As the stable homotopy groups $\tilde{\pi}_S^0(S^n)$ are finite, so are the groups $\tilde{\pi}_S^0(B_n G)$. Hence $\pi_S^0(BG) = \varprojlim_n \pi_S^0(B_n G)$, and $\pi_S^0(BG)$ is complete in the filtration topology.

It follows from the definition that $\alpha_G(I(G)) \subset [BG, Q_0 S^0] = F^1 \pi_S^0(BG)$, and since α_G is a ring homomorphism, $\alpha_G(I(G)^n) \subset F^n \pi_S^0(BG)$. Thus α_G is continuous and induces a homomorphism

$$\hat{\alpha}_G: \hat{A}(G) \rightarrow \pi_S^0(BG)$$

between the completions.

All the maps of 3.1 are continuous homomorphisms, when $R(G)$ is equipped with augmentation ideal topology and $K^*(BG)$ with a filtration topology similar to (F). Passing to completions we have

$$\begin{array}{ccc}
 \hat{A}(G) & \xrightarrow{\hat{\alpha}_G} & \pi_S^0(BG) \\
 \downarrow & & \downarrow e_* \\
 \hat{R}(G) & \xrightarrow{\hat{\alpha}} & K^*(BG)
 \end{array}$$

The main result of Atiyah [1] states that $\hat{\alpha}$ is an isomorphism. Therefore we can conclude the injectivity of $\hat{\alpha}_G$ if $\hat{A}(G)$ embeds into $\hat{R}(G)$. If G is cyclic then $A(G) \rightarrow R(G)$ is injective by Lemma 1.8. If moreover the order of G is a prime

power p^n , then the augmentation ideals of both rings have the p -adic topology by Proposition 1.12 and [4, p. 277]. But then $\hat{A}(G) \rightarrow \hat{R}(G)$ is injective, since the p -adic completion is an exact functor. We have proved

THEOREM 3.2. *Let G be a cyclic group of prime power order. Then $\hat{\alpha}_G$ is injective.*

We can express theorem 3.2 by saying that the maps $\alpha_G(x)$ for cyclic G are detected by K -theory. Indeed, the proof of 3.1 shows that $\alpha(x): BG \rightarrow BU \times \mathbb{Z}$ factors as

$$BG \xrightarrow{\alpha_G(x)} QS^0 \xrightarrow{e} BU \times \mathbb{Z} .$$

Since the map induced by $\alpha(x)$ in K -theory is non-trivial, if $x \neq 0$, so must be the one induced by $\alpha_G(x)$, too.

Next we invoke theorem 1.15 to show that the injectivity of $\hat{\alpha}_G$ can be deduced from that of $\hat{\alpha}_{G_p}$, for all Sylow subgroups G_p of G . Consider the commutative diagram

$$\begin{CD} \hat{A}(G) @>\hat{\alpha}_G>> \pi_S^0(BG) \\ @V \text{Res} VV @VVV \\ \bigoplus_p \hat{A}(G_p) @>\bigoplus \hat{\alpha}_{G_p}>> \bigoplus_p \pi_S^0(BG_p) \end{CD}$$

By Theorem 1.15, Res is injective. If the maps $\hat{\alpha}_{G_p}$ are injective for all $G_p \leq G$, then $\hat{\alpha}_G$ must be injective. Hence

THEOREM 3.3. *Let G be a finite group and $\{G_p\}$ its Sylow subgroups. If $\hat{\alpha}_{G_p}$ is injective for all $G_p \leq G$, then $\hat{\alpha}_G$ is injective.*

Theorem 3.3 reduces the study of $\hat{\alpha}_G$ to p -groups G . First, we note

LEMMA 3.4. *Let G be a p -group. Then $\hat{\alpha}_G$ is injective if and only if α_G is injective.*

(Indeed, as $A(G)$ embeds into $\hat{A}(G)$ by Corollary 1.11, one way is trivial and the converse follows since the p -adic completion is an exact functor and $I(G)$ and $\hat{\pi}_S^0(BG)$ both have p -adic topology, $I(G)$ by Proposition 1.12 and $\hat{\pi}_S^0(BG)$ being profinite with BG p -local [21].)

The smallest non-trivial p -group is the cyclic \mathbb{Z}/p , where we can apply Theorem 3.2. Suppose inductively that α_H is injective for all genuine subgroups

H of G . By naturality of α an element in $A(G)$ which has a non-zero restriction to some $H < G$, cannot lie in the kernel of α_G . Applying Theorem 1.3 we get

LEMMA 3.5. *Let G be a p -group. Suppose α_H is injective for all genuine subgroups $H < G$. Then α_G is injective on $\text{Ker } \chi_G$.*

To handle the rest, we have

PROPOSITION 3.6. *Let G be a p -group. There exists an element $x \in A(G)$ with $\chi_G(x) = p$ and $\chi_H(x) = 0$ for $H < G$. It is induced from an epimorphism $G \rightarrow (\mathbb{Z}/p)^d$.*

PROOF. The existence of x follows from the congruences of 1.3. However, we prefer to construct it directly.

Let $\Phi(G)$ be the Frattini subgroup of G , that is, the intersection of all maximal subgroups of G . We recall some elementary facts about $\Phi(G)$ [8, III § 3]:

- 1) $\Phi(G) = G^p[G, G]$,
- 2) $\Phi(G) \triangleleft G$ and $G/\Phi(G)$ is a maximal elementary abelian quotient of G , say $(\mathbb{Z}/p)^d$, and
- 3) the elements of $\Phi(G)$ are redundant in any set of generators for G .

One can also characterize the quotient $G/\Phi(G)$ as $H_1(G; \mathbb{Z}/p)$. In $A(\mathbb{Z}/p)^d$ we write down the element

$$y = p - \eta_1^{\text{tot}} + \eta_2^{\text{tot}} - \dots + (-1)^d p^{\binom{d-1}{2}} \eta_d$$

(See example 2.21, $y = pa_0$ in (2.22)). If $\pi: G \rightarrow G/\Phi(G)$ denotes the projection, then $\pi^*(y)$ has the required properties. Clearly $\chi_G(\pi^*(y)) = (\chi_{(\mathbb{Z}/p)^d}(y)) = p$. If $H < G$, then $\pi(H) < (\mathbb{Z}/p)^d$ and $\chi_H(\pi^*(y)) = \chi_{\pi(H)}(y) = 0$. Indeed, if $\pi(H) = (\mathbb{Z}/p)^d$, then $H\Phi(G) = G$, which implies $H = G$ by 3) above. This completes the proof of proposition 3.6.

LEMMA 3.7. $\mathbb{Z}x = \bigcap_{H < G} \text{Ker Res}_H^G$ and it is a λ -ideal of $A(G)$.

PROOF. The second claim follows from the first, since the maps Res_H^G are λ -homomorphisms. By definition $\text{Res}_H^G(x) = 0$ for each $H < G$. For the other containment suppose $\text{Res}_H^G(y) = 0$ for each $H < G$; we must show $\chi_G(y) \equiv 0 \pmod{p}$. Let $H < G$ be a subgroup of index p . As $H < N(H)$, H must be normal in G . The congruences of 1.3 become

$$0 = \chi_H(y) \equiv -\varphi(p)\chi_G(y) = -(p-1)\chi_G(y) \pmod{p}$$

thus $\chi_G(y) \equiv 0 \pmod{p}$.

We are ready to state the final result.

THEOREM 3.8. *Let G be a p -group. Suppose that*

- 1) α_H is injective for all $H < G$
- 2) α_G is injective for the λ -ideal Zx described in Proposition 3.6.

Then $\hat{\alpha}_G$ is injective.

PROOF. We first show that α_G is injective on $Zx \oplus \text{Ker } \chi_G$. If $m \in Z$, $\chi_G(y) = 0$ and $\alpha_G(mx + y) = 0$, then also

$$0 = \alpha_G(mx + y)\alpha_G(y) = \alpha_G(y^2)$$

since $xy = 0$ (all characters are 0). By Lemma 3.5 $y^2 = 0$, so $y = 0$. Thus $\alpha_G(mx) = 0$ and $m = 0$ by assumption 2).

Now any element in $A(G)$ can be written as $n + z$ with $0 \leq n < p$ and $z \in Zx \oplus \text{Ker } \chi_G$ since the latter ideal consists of z such that $\chi_G(z) \equiv 0 \pmod{p}$. Suppose $\alpha_G(n + z) = 0$, then

$$0 = \alpha_G(z)\alpha_G(n + z) = \alpha_G(nz + z^2)$$

where $nz + z^2 \in Zx \oplus \text{Ker } \chi_G$. From the above $nz = -z^2$, and taking characters we get

$$\chi_H(z) = 0 \text{ or } -n \text{ for all } H \leq G.$$

As $\chi_G(z) \equiv 0 \pmod{p}$ we must have $\chi_G(z) = 0$. But then $\chi_H(z) = 0$ for all $H < G$: if $H < G$ is a maximal subgroup with $\chi_H(z) = -n$ then by 1.3

$$-n = \chi_H(z) \equiv -\sum \varphi(K/H)\chi_K(z) = 0 \pmod{|N(H)/H|}$$

which is impossible, since $|N(H)/H|$ is a positive power of p . Thus $z = 0$ and $\alpha_G(n) = 0$ implies $n = \text{deg } \alpha_G(n) = 0$.

This completes the proof of Theorem 3.8.

4. Homological study of α_G .

In this section we shall study the maps induced by $\alpha_G(x): BG \rightarrow QS^0$ in homology for elementary abelian groups G . As a corollary we get that $\alpha_{(Z/p)^n}$ is injective. We obtain also information relative to the splitting $Q_0S_p^0 \cong J_p \times \text{cok } J_p$. We suppress the index G and write α for α_G .

Consider $\alpha(S)$ for a $(Z/p)^n$ -set S . The map α is additive, so we can restrict to transitive sets: $S = (Z/p)^n/H$ where $H \leq (Z/p)^n$. Both H and the quotient $(Z/p)^n/H = (Z/p)^n/H$ are elementary abelian, and S is induced from the regular representation $\eta_m = (Z/p)^m/1$ of $(Z/p)^m$. Thus $\alpha(S)$ factors

$$(4.1) \quad B(\mathbb{Z}/p)^n \rightarrow B(\mathbb{Z}/p)^m \xrightarrow{\alpha(\eta_m)} QS^0$$

Recall that the composition product in QS^0 corresponds to the product in $\coprod B\Sigma_n$ coming from the homomorphisms

$$\psi_{n,m}: \Sigma_n \times \Sigma_m \rightarrow \Sigma_{nm}$$

defined as

$$\psi_{n,m}(g, h)(i, j) = (g(i), h(j)).$$

Here Σ_{nm} is regarded as the permutation group of pairs (i, j) , $1 \leq i \leq n$, $1 \leq j \leq m$. This requires a linear ordering of the pairs; we use the lexicographic one.

We can express η_m inductively in terms of $\psi_{n,m}$. The first η_1 is just the inclusion $\mathbb{Z}/p \subset \Sigma_p$ as cyclic permutations. Then $\eta_2 = \psi_{p,p} \circ (\eta_1 \times \eta_1)$, and generally

$$\eta_n: (\mathbb{Z}/p)^n = \mathbb{Z}/p \times (\mathbb{Z}/p)^{n-1} \xrightarrow{\eta_1 \times \eta_{n-1}} \Sigma_p \times \Sigma_{p^{n-1}} \xrightarrow{\psi_{p,p^{n-1}}} \Sigma_{p^n}.$$

Hence $\alpha(\eta_n)$ can be written as the composition

$$(4.2) \quad \alpha(\eta_n): B(\mathbb{Z}/p)^n = (B\mathbb{Z}/p)^n \xrightarrow{(B\eta_1)^n} (B\Sigma_p)^n \xrightarrow{i^n} (QS^0)^n \xrightarrow{\cdot} QS^0.$$

We shall need certain facts about the homology of QS^0 with \mathbb{Z}/p -coefficients. General references for this are [10] and [12] for $p=2$ and [6] for $p>2$. Here is a summary.

The space QS^0 has two products: the loop sum $*$ and the composition product \cdot . They induce products on $H_*(QS^0; \mathbb{Z}/p)$, denoted similarly. They are homomorphisms $Q^b: H_*(QS^0; \mathbb{Z}/p) \rightarrow H_*(QS^0; \mathbb{Z}/p)$ with the following properties [modifications for the case $p=2$ are stated inside square brackets]:

$$(4.3) \quad \text{Degree:} \quad Q^b \text{ raises degree by } 2b(p-1) \quad [b]$$

$$(4.4) \quad \text{Evaluation:} \quad \begin{array}{ll} Q^b x = 0 & \text{if } 2b < \deg x \quad [b < \deg x] \\ Q^b x = x^{*p} & \text{if } 2b = \deg x \quad [b = \deg x] \end{array}$$

$$(4.5) \quad \text{Cartan formula: } Q^b(x * y) = \sum_{i+j=b} Q^i x * Q^j y.$$

$$(4.6) \quad \text{Adem relations: If } a > pb \text{ then}$$

$$Q^a Q^b x = \sum (-1)^{a+i} \binom{(p-1)(t-b)-1}{pt-a} Q^{a+b-i} Q^i x;$$

if $p > 2$, $a \geq pb$ and β denotes the mod p Bockstein, then

$$Q^a \beta Q^b x = \sum (-1)^{a+i} \binom{(p-1)(t-b)}{pt-a} \beta Q^{a+b-i} Q^i x$$

$$+ \sum (-1)^{a+t} \binom{(p-1)(t-b)-1}{pt-a-1} Q^{a+b-t} \beta Q^t x.$$

In all cases the summation is over t such that $(p+1)t \geq a+b$.

(4.7) Nishida relations: If P_*^r is dual to the reduced p th power P^r [the square Sq^r] then

$$P_*^a Q^b x = \sum_{t \geq 0} (-1)^{a+t} \binom{(p-1)(b-a)}{a-pt} Q^{b-a+t} P_*^t x;$$

if $p > 2$ then

$$\begin{aligned} P_*^a \beta Q^b x &= \sum_{t \geq 0} (-1)^{a+t} \binom{(p-1)(b-a)-1}{a-pt} \beta Q^{b-a+t} P_*^t x \\ &+ \sum_{t \geq 0} (-1)^{a+t} \binom{(p-1)(b-a)-1}{a-pt-1} Q^{b-a+t} P_*^t \beta x. \end{aligned}$$

Let $[k] \in H_0(QS^0; \mathbb{Z}/p)$ denote the component of maps of degree k . E. Dyer and R. Lashof showed that the homology ring $H_*(QS^0; \mathbb{Z}/p)$ was generated by successive operations of $Q^a, \beta Q^a$ on $[1]$ as an algebra under $*$. To make a precise statement, we introduce the Dyer–Lashof algebra $R(p)$.

Let \mathcal{F} be the free graded associative algebra generated by the symbols $Q^s, s \geq 0$ and $\beta Q^s, s > 0$ with degrees $2s(p-1)$ and $2s(p-1)-1$ respectively [if $p=2$, \mathcal{F} is generated by $Q^s, s \geq 0$, with degree s]. The monomials in \mathcal{F} can be written as

$$\beta^{\varepsilon_1} Q^{s_1} \dots \beta^{\varepsilon_k} Q^{s_k}$$

with $\varepsilon_i = 0$ or 1 and $s_i \geq \varepsilon_i$. Denote them by Q^I , where $I = (\varepsilon_1, s_1, \dots, \varepsilon_k, s_k)$. We say that I is admissible if $s_1 \leq ps_2 - \varepsilon_2, \dots, s_{k-1} \leq ps_k - \varepsilon_k$, and we define the length and excess of I by $l(I) = k$ and

$$\begin{aligned} e(I) &= (2s_1 - \varepsilon_1) - \sum_{j=2}^k (2s_j(p-1) - \varepsilon_j) \quad (p > 2) \\ e(I) &= s_1 - \sum_{j=2}^k s_j \quad (p = 2) \end{aligned}$$

The quotient of \mathcal{F} by the ideal generated by the Adem relations and by monomials with $e(I) < 0$ is the Dyer–Lashof algebra $R(p)$.

The formulas (4.3)–(4.6) tell that $R(p)$ acts on $H_*(QS^0; \mathbb{Z}/p)$. In fact the set

$$(4.8) \quad X = \{Q^I[1], I \text{ admissible}, e(I) + \varepsilon_1 > 0\}$$

forms a basis for the $*$ -algebra $H_*(QS^0; \mathbb{Z}/p)$ up to component shift. Indeed, let $\mathbb{Z}/p[\mathbb{Z}]$ be the group ring of $\mathbb{Z} = \pi_0(QS^0)$. Then

$$H_*(QS^0; \mathbf{Z}/2) = PX \otimes \mathbf{Z}/2[\mathbf{Z}]$$

$$H_*(QS^0; \mathbf{Z}/p) = PX^+ \otimes EX^- \otimes \mathbf{Z}/p[\mathbf{Z}], \quad \text{if } p > 2,$$

where P and E denote the polynomial and exterior algebras, respectively, and X^+ (X^-) is the even (odd) degree part of X .

The composition product is related to the operations Q^b by May's formula:

$$(4.9) \quad \begin{aligned} Q^b(x) \cdot f &= \sum_{t \geq 0} Q^{b+t}(x \cdot P_*^t f) \quad \text{and, if } p > 2 \\ \beta Q^b(x) \cdot f &= \sum \beta Q^{b+t}(x \cdot P_*^t f) - (-1)^{\deg x} \sum Q^{b+t}(x \cdot P_*^t \beta f) \end{aligned}$$

After these preparations we turn to the evaluation of (4.2) in homology. The map $i: B\Sigma_p \rightarrow QS^0$ is obtained from the Dyer–Lashof map θ_p as the composite

$$i: B\Sigma_p = E\Sigma_p \times_{\Sigma_p} (*)^p \rightarrow E\Sigma_p \times_{\Sigma_p} (QS^0)^p \xrightarrow{\theta_p} QS^0,$$

where $*$ goes to the identity map in the 1-component Q_1S^0 . By (4.2), $\alpha(\eta_1)$ is of the form $B\mathbf{Z}/p \xrightarrow{B\eta_1} B\Sigma_p \rightarrow QS^0$. This is precisely the map used in the definition of Q^s [6, pp. 7–8]; if $e_m \in H_m(B\mathbf{Z}/p; \mathbf{Z}/p)$ denotes the standard generator then

$$(4.10) \quad \alpha(\eta_1)_*(e_m) = Q_m[1] = \begin{cases} (-1)^s Q^s[1] & \text{if } m = 2s(p-1) \\ (-1)^s \beta Q^s[1], & \text{if } m = 2s(p-1) - 1 \text{ for } p \text{ odd,} \\ 0 & \text{otherwise} \end{cases}$$

$$= Q^m[1] \quad \text{for } p=2.$$

It follows from (4.2) and (4.10) that $\alpha(\eta_n)_*$ takes the generators $e_{i_1} \otimes \dots \otimes e_{i_n}$ of $H_*(B(\mathbf{Z}/p)^n; \mathbf{Z}/p)$ to products of the form

$$\pm \beta^{s_1} Q^{s_1}[1] \cdot \dots \cdot \beta^{s_n} Q^{s_n}[1].$$

We would like to express these elements in the $*$ -product basis (4.8). A two-fold product, for example $Q^a[1] \cdot Q^b[1]$, becomes

$$(4.11) \quad \begin{aligned} Q^a[1] \cdot Q^b[1] &= \sum_{t \geq 0} Q^{a+t}(P_*^t Q^b[1]) \\ &= \sum_{t \geq 0} (-1)^t \binom{(p-1)(b-t)}{t} Q^{a+t} Q^{b-t}[1] \end{aligned}$$

by (4.9) and (4.7). Applying the Adem relations (4.6) it can be written as a linear combination of admissible terms $Q^s Q^t[1]$. If the excess is negative then $Q^t[1] = 0$ by (4.4). Similarly it is shown by induction on the length of the product that

LEMMA 4.12.

$$\beta^{\varepsilon_1} Q^{\varepsilon_1}[1] \cdot \dots \cdot \beta^{\varepsilon_n} Q^{\varepsilon_n}[1] = \sum \lambda_I Q^I[1]$$

where I ranges over admissible sequence of length n and excess ≥ 0 .

The terms $Q^I[1]$ with $e(I) + \varepsilon_1 = 0$ decompose as $*$ -products of shorter $Q^J[1]$'s (4.4).

We shall now find the special case of Lemma 4.12 in the lowest degree where we can get an admissible $Q^I[1]$ of length n and excess > 0 involving no Bocksteins β . It is clearly $Q^{p^n} Q^{p^{n-1}} \dots Q^1[1]$ in dimension $2(p^{n+1} - 1)$ with excess 2 [if $p=2$ then $d=2^{n+1} - 1$ and $e=1$]. We give the proofs of the next two lemmas only for $p > 2$. The (easier) case $p=2$ follows by trivial modifications.

LEMMA 4.13. $Q^{p^n}[1] \cdot Q^{p^{n-1}}[1] \cdot \dots \cdot Q^1[1] = Q^{p^n} Q^{p^{n-1}} \dots Q^1[1]$.

PROOF. To begin with, $Q^p[1] \cdot Q^1[1] = Q^p Q^1[1]$ by (4.11). Suppose by induction that the claim holds for n . Since $x_n = Q^{p^n} Q^{p^{n-1}} \dots Q^1[1]$ is primitive, so is also $P_*^t x_n$. If $t > 0$, then according to (4.7) $P_*^t x_n$ is a linear combination of $Q^I[1]$'s of length n and degree $< 2(p^{n+1} - 1)$, without Bocksteins. By the minimality of x_n , $P_*^t x_n$ is $*$ -decomposable. By a general theorem of Hopf algebras [15, Proposition 4.23] $P_*^t x_n$ must then be a $*$ - p th power, especially

$$\deg P_*^t x_n = 2(p^{n+1} - 1) - 2t(p - 1) \equiv -2 + 2t \equiv 0 \pmod{p}$$

so that $t \equiv 1 \pmod{p}$. Now we can apply May's formula (4.9) to get

$$\begin{aligned} Q^{p^{n+1}}[1] \cdot Q^{p^n}[1] \cdot \dots \cdot Q^1[1] &= Q^{p^{n+1}}[1] \cdot x_n \\ &= \sum_{t \geq 0} Q^{p^{n+1} + t} (P_*^t x_n) = Q^{p^{n+1}} x_n = Q^{p^{n+1}} Q^{p^n} \dots Q^1[1] \end{aligned}$$

since $Q^s(x^{*p}) \neq 0$ only if $s \equiv 0 \pmod{p}$ in virtue of the Cartan formula (4.5).

Now we can prove that the map $\alpha: A((\mathbb{Z}/p)^n) \rightarrow \pi_3^0(B(\mathbb{Z}/p)^n)$ is injective on $Z\eta_n$ and thereby on the whole of $A((\mathbb{Z}/p)^n)$.

PROPOSITION 4.14. $\alpha(m\eta_n)$ is homologically non-trivial for all non-zero integers m .

PROOF. Let first $m > 0$. Then using the diagonal formula

$$\psi(e_{2i}) = \sum e_{2i_1} \otimes \dots \otimes e_{2i_m}, \quad i_1 + \dots + i_m = i$$

for $B\mathbb{Z}/p \rightarrow (B\mathbb{Z}/p)^m$ and Lemmas 4.12 and 4.13 we obtain

$$\alpha(m\eta_n)_*(e_{2mp^{n-1}} \otimes \dots \otimes e_{2m(p-1)}) = (Q^{p^n} Q^{p^{n-1}} \dots Q^1[1])^{*m} + \dots$$

where the other terms are of the form $Q^{l_1}[1]^* \dots * Q^{l_n}[1]$ with $l(I_j) = n$ and $\deg(I_j) < 2(p^{n+1} - 1)$ for at least one j . Since $Q^{p^n} Q^{p^{n-1}} \dots Q^1[1]$ is a polynomial generator, they cannot cancel the first term.

If $m < 0$, then apply the loop inverse χ_* , and note that $\chi_*(x) = x * [-2 \deg x]$ on primitive elements x .

THEOREM 4.15. $\hat{\alpha}: \hat{A}((\mathbb{Z}/p)^n) \rightarrow \pi_s^0 \mathfrak{p}(B(\mathbb{Z}/p)^n)$ is injective for all primes p .

PROOF. By Theorems 3.2 and 3.8 we are reduced to showing that α is injective on $\mathbb{Z}x$, where

$$x = p - \eta_1^{\text{tot}} + \dots + (-1)^n p^{\binom{n-1}{2}} \eta_n.$$

The argument of Proposition 4.14 applies also here, since the terms η_i^{tot} contribute in homology only by $*$ -products of $Q^l[1]$ with $l(I) < n$ (cf. (4.1)).

This completes the proof of theorem 4.15.

Let $Q_0 S^0$ be the 0-component of $Q S^0$. Let X_p denote the localization of the space X at a prime p [21]. D. Sullivan has showed that the space $Q_0 S^0$ splits locally

$$(4.16) \quad Q_0 S_p^0 \cong J_p \times \text{cok } J_p.$$

The space J_2 is defined as the fibre of $\psi^3 - 1: BO_2 \rightarrow B\text{Spin}_2$. At odd primes J_p is the fibre of $\psi^k - 1: BU_p \rightarrow BU_p$, where k is a prime power generating the group of units in \mathbb{Z}/p^2 . The homotopy groups of J_p are essentially the p -primary part of the image of the J -homomorphism $O \rightarrow G$ in the stable homotopy of spheres. To describe the second factor $\text{cok } J_p$ we recall the discrete models for J_p due to D. Quillen [14, chapter VIII].

First, let $p=2$. Let F_3 denote the finite field with 3 elements. Let $N_n(F_3)$ be the group of orthogonal transformations of the quadratic space $(F_3^n, x_1^2 + \dots + x_n^2)$ for which the determinant and the spinor norm [14, p. 164] agree. We encounter now a similar situation to the construction of $Q S^0$ from the symmetric groups: there are sum and product maps on the disjoint union

$$(4.17) \quad \coprod_{n \geq 0} BN_n(F_3)$$

coming from direct sum and tensor product of quadratic spaces.

Let \bar{F}_3 be an algebraic closure of F_3 and choose an embedding $\mu: \bar{F}_3^* \rightarrow \mathbb{C}^*$. If G is a finite group and $\varrho: G \rightarrow \text{Gl}_n(\bar{F}_3)$ a representation of G , then the complex-valued function on G

$$\chi(g) = \sum_{i=1}^n \mu(\lambda_i)$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of $\varrho(g)$, is the character of a unique element in the complex representation ring $R(G)$. Moreover, Quillen has proved that if ϱ takes values in $O_n(\bar{F}_3)$, then χ is the character of an element in the real representation ring $RO(G)$.

We lift the standard representations of $N_n(F_3)$ in \bar{F}_3^n in the above way to virtual representations in $RO(N_n(F_3))$ and apply $\alpha: RO(G) \rightarrow KO(BG)$ to get maps

$$v_n: BN_n(F_3) \rightarrow BO \times (n).$$

They are compatible with the sum and product on (4.17), giving rise to an H -map

$$v: \Omega B \left(\prod_{n \geq 0} BN_n(F_3) \right) \rightarrow BO \times \mathbb{Z}$$

from its group completion. Now the Adams operation ψ^3 is characterized by its action on the characters $(\psi^3 \chi)(g) = \chi(g^3)$, so $\psi^3 \circ v_n = v_n$ as the Frobenius map $\lambda \rightarrow \lambda^3$ just permutes the eigenvalues of any representation realizable over F_3 .

Let J_2^0 denote the zero component of $\Omega B \left(\prod_{n \geq 0} BN_n(F_3) \right)$ localized at 2. Then $v: J_2^0 \rightarrow BO_2$ lifts to an H -map $J_2^0 \rightarrow J_2$, which can be proved to be a homotopy equivalence e.g. by cohomological methods. From now on we identify J_2^0 and J_2 .

Let

$$e: QS^0 \cong \Omega B \left(\prod_{n \geq 0} B\Sigma_n \right) \rightarrow \Omega B \left(\prod_{n \geq 0} BN_n(F_3) \right)$$

be induced from the functor which takes a finite set S to the vector space F_3^S . We restrict e to the zero component and localize to get $e: Q_0S_2^0 \rightarrow J_2$. We shall also use the analogous map $e: Q_0S_2^0 \rightarrow BO_2$, induced from the functor $S \mapsto \mathbb{R}^S$. Then the triangle

(4.18) 

clearly commutes. The space $\text{cok } J_2$ is defined as the fibre in

$$(4.19) \quad \text{cok } J_2 \rightarrow Q_0S_2^0 \xrightarrow{e} J_2.$$

There exists a splitting $\alpha_2: J_2 \rightarrow Q_1S_2^0$, and $\alpha_2 * [-1]$ gives (4.16).

At odd primes p the model for J_p is constructed from general linear groups

over the finite field $F_k: J_p$ is equivalent to the zero component of the group completion of

$$(4.17) \quad \coprod_{n \geq 0} BGL_n(F_k), \quad k \text{ a prime power generating } (\mathbb{Z}/p^2)^* .$$

As before defines $e: Q_0S_p^0 \rightarrow J_p$ and gets a commutative diagram

$$(4.18') \quad \begin{array}{ccc} & & J_p \\ & \nearrow e & \downarrow \\ Q_0S_p^0 & & BU_p \\ & \searrow e & \end{array}$$

and $\text{cok } J_p$ is defined as the fibre in

$$(4.19') \quad \text{cok } J_p \rightarrow Q_0S_p^0 \xrightarrow{e} J_p .$$

Let now G be a p -group. We shall in the following consider the abelian groups $[BG, X_p]$, where $X = Q_0S^0, BU, U$ for all p and in addition to these, $X = BO$ and SO for $p=2$. We claim that in all cases

$$[BG, X_p] = [BG, X] .$$

If $X = Q_0S^0$ this holds because Q_0S^0 has finite homotopy groups, so $Q_0S^0 = \prod_p Q_0S_p^0$, and BG is p -local (even p -complete) [21, section 3].

For the other spaces we recall the results of Atiyah [1] and Atiyah–Segal [2]. Consider the representable K -theory and the theory $K^*(; \mathbb{Z}_{(p)})$ defined by the unitary spectrum and its localization at p . For any finite CW-complex Y we have

$$K^*(Y; \mathbb{Z}_{(p)}) \cong K^*(Y) \otimes \mathbb{Z}_{(p)} .$$

The formula is valid also for $Y = BG$ since it follows from [1] and [2] that \lim^1 of the inverse systems $K^*(B_nG)$ and $K^*(B_nG) \otimes \mathbb{Z}_{(p)}$ vanishes, so that $K^*(\overleftarrow{BG}) = \lim K^*(B_nG)$ and $K^*(BG; \mathbb{Z}_{(p)}) = \lim K^*(B_nG) \otimes \mathbb{Z}_{(p)}$. For any group G $K^0(\overleftarrow{BG}) = \hat{R}(G)$ and $K^1(BG) = 0$ [1, p. 270] and in the case of p -groups the completion is the p -adic one: $\tilde{K}^0(BG) = I(G) \otimes \hat{\mathbb{Z}}_p$ [4 p. 277]. Since these groups are clearly unaffected by $\otimes \mathbb{Z}_{(p)}$, we get

$$(4.20) \quad [BG, BU] = [BG, BU_p] = I(G) \otimes \hat{\mathbb{Z}}_p, \quad [BG, U_p] = 0$$

where $I(G)$ is the augmentation ideal of $R(G)$.

If $p=2$, then using the Real K -theory KR^* instead of K^* and [2, p. 17] we obtain in the same fashion

$$(4.21) \quad [BG, BO] = [BG, BO_2] = I(G) \otimes \hat{\mathbb{Z}}_2 ,$$

$$[BG, SO_2] = \text{vector space over } \mathbb{Z}/2$$

for 2-groups G , where $I(G)$ is the augmentation ideal of $R_{\mathbb{R}}(G)$.

Let G be a p -group. After these preliminaries we turn to the question: when does a map $\hat{\alpha}(x): BG \rightarrow Q_0S^0$ lift to $\text{cok } J_p$ in the fibration (4.19), (4.19'). In order for $e \circ \hat{\alpha}(x): BG \rightarrow Q_0S^0 \rightarrow J_p$ to be nulhomotopic, it is necessary in the light of (4.18) and (4.18') that the image of $\hat{\alpha}(x)$ under $e_*: \tilde{\pi}_s^0(BG) \rightarrow \tilde{K}^0(BG)$ is zero. From Proposition 3.1, this is equivalent to

$$x \in \hat{A}_0(G) = \text{Ker} (\hat{A}(G) \rightarrow \hat{R}(G)).$$

If p is odd, this condition is also sufficient, since in the mapping sequence of the fibration $J_p \rightarrow BU_p \xrightarrow{\psi^p-1} BU_p$

$$[BG, U_p] \rightarrow [BG, J_p] \rightarrow [BG, BU_p]$$

the first group is trivial (4.20), so $x \in \hat{A}_0(G)$ maps to zero already in $[BG, J_p]$.

In particular, if G is an elementary abelian group $(\mathbb{Z}/p)^n$ with odd p , we know that all the maps $\hat{\alpha}(x): BG \rightarrow Q_0S^0$, $x \in \hat{A}_0(G)$ are homotopically distinct (Theorem 4.15). Thus $\hat{\alpha}$ lifts to a monomorphism $\hat{\alpha}'$

$$\begin{array}{ccc} & \hat{A}_0(G) & \\ & \swarrow \hat{\alpha}' & \downarrow \hat{\alpha} \\ [BG, \text{cok } J_p] & \rightarrow & [BG, Q_0S^0] \rightarrow [BG, J_p]. \end{array}$$

THEOREM 4.22. Let p be an odd prime and G the elementary abelian group $(\mathbb{Z}/p)^n$. Then the ideal

$$\hat{A}_0(G) = \text{Ker} (\hat{A}(G) \rightarrow \hat{R}(G))$$

maps monomorphically into $[BG, \text{cok } J_p]$.

Let then G be a 2-group and $x \in \hat{A}_0(G)$. Then the image of $\hat{\alpha}(x)$ is 0 in $[BG, BU_2]$. To see when $e \circ \hat{\alpha}(x): BG \rightarrow J_2$ is non-trivial, we consider the maps $J_2 \rightarrow BO_2 \xrightarrow{c} BU_2$, where c is complexification. The map $[BG, BO_2] \rightarrow [BG, BU_2]$ corresponds to the completion of $R_{\mathbb{R}}(G) \subset R(G)$ by (4.20), (4.21) and [2, p. 17], which is injective. In the mapping sequence of $J_2 \rightarrow BO_2 \xrightarrow{\psi^2-1} B\text{Spin}_2$

$$[BG, \text{Spin}_2] \rightarrow [BG, J_2] \rightarrow [BG, BO_2]$$

the first group is a subgroup of $[BG, SO_2]$ as $SO_2 \cong \mathbb{R}P^\infty \times \text{Spin}_2$, hence it is a vector space over $\mathbb{Z}/2$. Thus (at least) $2x$ maps to zero in $[BG, J_2]$. We have proved the first half of

THEOREM 4.23. *Let G be the elementary abelian 2-group $(\mathbb{Z}/2)^n$. Then the ideal $2\hat{A}_0(G)$ maps monomorphically into $[BG, \text{cok } J_2]$. There are elements in $\hat{A}_0(G)$ which do not lift to $\text{cok } J_2$.*

PROOF. Consider the critical element $x \in A_0(G)$ with $\chi_G(x)=2$ and $\chi_H(x)=0$ for all genuine subgroups $H < G$. We claim that $1-x$ can be written as a product in terms of the 2^n-1 quotients $\eta_1^i = G/(\mathbb{Z}/2)^{n-1}$:

$$1-x = \prod_{i=1}^{2^n-1} (\eta_1^i - 1).$$

Indeed, check the characters. First $\chi_H(\eta_1^i)=2$ or 0 according to whether $H \leq (\mathbb{Z}/2)^{n-1}$ or $H \not\leq (\mathbb{Z}/2)^{n-1}$, the hyperplane defining η_1^i . Therefore we get

$$\chi_G\left(\prod_{i=1}^{2^n-1} (\eta_1^i - 1)\right) = (-1)^{2^n-1} = -1.$$

On the other hand each hyperplane containing H corresponds to a line inside H^\perp . If $H < G$ the number of these, $|H|^\perp - 1$, is odd, so

$$\chi_H\left(\prod_{i=1}^{2^n-1} (\eta_1^i - 1)\right) = 1^{\text{odd}}(-1)^{\text{even}} = 1, \quad H < G.$$

Thus $\alpha(1-x)$ is a composition product of maps of the form

$$BG \xrightarrow{B\eta_1^i} B\mathbb{Z}/2 \xrightarrow{i_2} Q_2S^0 \xrightarrow{*[-1]} SG.$$

But the map $i_2 * [-1]: B\mathbb{Z}/2 \rightarrow SG$ is homotopy equivalent to the composite

$$RP^\infty \rightarrow SO \xrightarrow{J} SG$$

[6, p. 120] so that $\alpha(1-x)$ factors through $J: SO \rightarrow SG$. Let $e_1: SG \rightarrow J^\otimes$ be the 1-component of the map e defined just before (4.18). We showed in 4.14 that $\alpha(-x)$, hence $\alpha(1-x)$ induces a non-trivial map in homology. It is well-known that the composite

$$H_*(SO) \xrightarrow{J_*} H_*(SG) \xrightarrow{e_{1*}} H_*(J)$$

is injective [6, p. 120 and Theorem 12.5 p. 185]. Then $e_1 \circ \alpha(1-x)$, hence $e \circ \alpha(-x)$ must be homologically non-trivial.

This completes the proof of Theorem 4.23.

REMARK 4.24. Theorems 4.22 and 4.23 enable us to get hold of elements in $H_*(\text{cok } J_p; \mathbb{Z}/p)$. Let us consider the first case $A_0(\mathbb{Z}/2 \oplus \mathbb{Z}/2) = \mathbb{Z}x$ ($A_0(\mathbb{Z}/2) = 0!$). It is most convenient to evaluate $f = \alpha(1-x)$, since from the preceding proof

$$1-x = (\eta_1^1 - 1)(\eta_1^2 - 1)(\eta_1^3 - 1)$$

where $\eta_1^i: \mathbf{Z}/2 \oplus \mathbf{Z}/2 \rightarrow \mathbf{Z}/2$ are projections to the first and second factor for $i = 1, 2$, and η_1^3 takes the quotient modulo the diagonal subgroup $\Delta \mathbf{Z}/2 \subset \mathbf{Z}/2 \oplus \mathbf{Z}/2$.

The maps $f_i = \alpha(\eta_1^i - 1): \mathbf{RP}^\infty \times \mathbf{RP}^\infty \rightarrow \mathbf{RP}^\infty$, $i = 1, 2, 3$, have the effect

$$f_{1*}(e_m \otimes e_n) = \delta_{n0} x_m, f_{2*}(e_m \otimes e_n) = \delta_{m0} x_n$$

and

$$f_{3*}(e_m \otimes e_n) = \binom{m+n}{m} x_{m+n}$$

on homology (cf. 4.10). Here $x_k = Q^k[1] * [-1] \in H_*(SG; \mathbf{Z}/2)$, and adding up we get

$$(4.25) \quad f_*(e_m \otimes e_n) = \sum_{i=0}^m \sum_{j=0}^n \binom{m+n-i-j}{m-i} x_i \cdot x_j \cdot x_{m+n-i-j}.$$

As a special case of this formula $f_*(e_{2n} \otimes e_1) = p_{2n+1}$, where the polynomial

$$p_{2n+1} = x_{2n+1} + \sum_{i=1}^n x_i x_{2n+1-i}$$

is the standard primitive element of degree $2n+1$ in the subalgebra $E(x_1, x_2, \dots) \subset H_*(SG)$.

Let \bar{f} denote $\alpha(-x) = f * [-1]$. We know from theorem 4.23 that $2\bar{f}: (\mathbf{RP}^\infty)^2 \rightarrow Q_0 S^0$ lifts to $\text{cok } J_2$. Therefore the elements

$$\begin{aligned} C_{4n+2} &= (2\bar{f})_*(e_{4n} \otimes e_2) = \bar{f}_*(e_{2n} \otimes e_1) * \bar{f}_*(e_{2n} \otimes e_1) \\ &= p_{2n+1} * p_{2n+1} * [-2] \end{aligned}$$

$n \geq 1$, lie in $\text{Ker } e_*$. Since they are primitive, they lie in $H_*(\text{cok } J_2, \mathbf{Z}/2)$. (The elements C_{2^i-2} have a connection with the Arf invariant conjecture: they are spherical if and only if there are stable homotopy classes in $\pi_{2^i-2}^S(S^0)$ of Arf invariant one [9].)

REMARK 4.26. We succeeded in proving that $\hat{\alpha}_G$ is injective for elementary abelian G by evaluating the maps $\alpha_G(nx)$ in homology. Let us indicate briefly where this program fails for more complicated groups. The smallest ones we have not covered are the following three groups of order 8:

$$\begin{aligned} \mathbf{Z}/4 \oplus \mathbf{Z}/2 &= \langle x, y \mid x^4 = y^2 = 1, xy = yx \rangle \\ \text{D8} &= \langle x, y \mid x^4 = y^2 = 1, y^{-1}xy = x^3 \rangle \\ \text{Q8} &= \langle x, y \mid x^4 = 1, y^2 = x^2, y^{-1}xy = x^3 \rangle \end{aligned}$$

(D8 and Q8 are the dihedral and the quaternion groups). In all cases the Frattini subgroup $\Phi(G)=G^2$ is $\mathbb{Z}/2$ generated by x^2 . The cohomology of G (with $\mathbb{Z}/2$ coefficients) can be computed from the spectral sequence of the central extension

$$1 \rightarrow \Phi(G) \rightarrow G \xrightarrow{\pi} \mathbb{Z}/2 \oplus \mathbb{Z}/2 \rightarrow 1.$$

The E^2 -term is

$$H^*(\mathbb{Z}/2) \otimes H^*(\mathbb{Z}/2 \oplus \mathbb{Z}/2) = P(t) \otimes P(t_1, t_2).$$

We choose t_1 and t_2 as the generators of the cohomology of $\langle \pi(x) \rangle$ and $\langle \pi(y) \rangle$. The differentials are determined by the characteristic class $d_2(t) \in H^2(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$, which is

$$t_1^2, t_1^2 + t_1 t_2 \quad \text{and} \quad t_1^2 + t_1 t_2 + t_2^2,$$

respectively.

The critical elements $BG \rightarrow QS^0$ are compositions of

$$BG \xrightarrow{B\pi} B(\mathbb{Z}/2 \oplus \mathbb{Z}/2) = (\mathbb{R}P^\infty)^2$$

with the maps $\alpha(nx): (\mathbb{R}P^\infty)^2 \rightarrow QS^0$. We evaluated $\bar{f} = \alpha(-x)$ in the preceding remark. From (4.25) we get

$$(4.27) \quad \bar{f}_*(e_n \otimes e_m) = \bar{f}_*(e_m \otimes e_n), \quad \bar{f}_*(e_n \otimes e_0) = 0 \quad (n > 0).$$

Consider now e.g. the cohomology of $G = D8$. In its spectral sequence

$$d_3(t^2) = d_3(\text{Sq}^1 t) = \text{Sq}^1 d_2 t = \text{Sq}^1 (t_1^2 + t_1 t_2) = t_1^2 t_2 + t_1 t_2^2 = t_2 d_2 t = 0$$

so that $E^3 = E^\infty$ and

$$H^*(D8) = P(s) \otimes P(t_1, t_2) / (t_1^2 + t_1 t_2)$$

where $s \in H^2(D8)$ is any element whose image is $t^2 \in H^2(\mathbb{Z}/2)$, and t_1 and t_2 come from $H^*(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$. Thus the image of $B\pi^*$ in $H^n(D8)$ is generated by the elements $t_1^n = t_1^{n-1} t_2 = \dots = t_1 t_2^{n-1}$ and t_2^n . Dually the image of $B\pi_*$ in $H_n(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$ is generated by

$$e_n \otimes e_0 + e_{n-1} \otimes e_1 + \dots + e_1 \otimes e_{n-1} \quad \text{and} \quad e_0 \otimes e_n.$$

From (4.27) $\bar{f}_*(\text{Im } B\pi_*) = 0$. Hence all maps $\alpha_{D8}(nx) = (-n\bar{f}) \circ B\pi$ vanish in $\mathbb{Z}/2$ -homology.

A similar computation shows that $\bar{f} \circ B\pi$ induces the zero map $H_*(Q8) \rightarrow H_*(Q_0 S^0)$. In fact here $\text{Im } B\pi^* = 0$ from dimension 4 on. Finally for $G = \mathbb{Z}/4 \oplus \mathbb{Z}/2$ we get that $(\bar{f} \circ B\pi)_*$ is non-trivial precisely in dimension 3. But then $(2\bar{f} \circ B\pi)_*$ vanishes.

By Proposition 3.1 these maps induce 0 also in K -theory. We pose the

QUESTION. Are the maps

$$f_n: BG \xrightarrow{B\pi} \mathbb{R}P^\infty \times \mathbb{R}P^\infty \xrightarrow{\alpha(n,x)} Q_0S^0,$$

where $G = \mathbb{Z}/4 \oplus \mathbb{Z}/2$, $D8$, $Q8$ and $x = 2 - \eta_1^1 - \eta_1^2 - \eta_1^3 + \eta_2 \in A_0(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$ homotopic to zero?

REFERENCES

1. M. F. Atiyah, *Characters and cohomology of finite groups*, Inst. Hautes Études Sci. Publ. Math. 9 (1961), 247–288.
2. M. F. Atiyah and G. B. Segal, *Equivariant K-theory and completion*, J. Differential Geometry 3 (1969), 1–18.
3. M. F. Atiyah and G. B. Segal, *Exponential isomorphisms for λ -rings*, Quart. J. Math. Oxford Ser. 22 (1971), 371–378.
4. M. F. Atiyah and D. O. Tall, *Group representations, λ -rings and the J-homomorphism*, Topology 8 (1969), 253–297.
5. W. Burnside, *Theory of groups of finite order*, 2nd Edition 1911. Reprinted by Dover 1955.
6. F. R. Cohen, T. J. Lada and J. P. May, *The homology of iterated loop spaces*, Lecture Notes in Mathematics 533, Springer-Verlag, Berlin - Heidelberg - New York, 1976.
7. A. Dress, *A characterisation of solvable groups*, Math. Z. 110 (1969), 213–217.
8. B. Huppert, *Endliche Gruppen I*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Band 134, Springer-Verlag, Berlin - Heidelberg - New York, 1967.
9. I. Madsen, *On the action of the Dyer–Lashof algebra in $H_*(G)$ and $H_*(G/TOP)$* , Thesis, University of Chicago, 1970.
10. I. Madsen, *On the action of the Dyer–Lashof algebra in $H_*(G)$* , Pacific J. Math. 60 (1975), 235–275.
11. I. Madsen, *Smooth spherical space forms in Geometric Applications of Homotopy Theory I*, Proceedings, Evanston, 1977, Lecture Notes in Mathematics 657, Springer-Verlag, Berlin - Heidelberg - New York, 1978.
12. I. Madsen and R. J. Milgram, *The cobordism of topological and PL-manifolds*, to appear in Annals of Mathematics Studies, Princeton University Press, Princeton.
13. J. P. May, *The geometry of iterated loop spaces*, Lectures Notes in Mathematics 271, Springer-Verlag, Berlin - Heidelberg - New York, 1972.
14. J. P. May (with contributions by F. Quinn, N. Ray and J. Tornehave), *E_∞ ring spaces and E_∞ ring spectra*, Lecture Notes in Mathematics 577, Springer-Verlag, Berlin - Heidelberg - New York, 1977.
15. J. W. Milnor and J. C. Moore, *On the structure of the Hopf algebras*, Ann. of Math. 81 (1965), 211–264.
16. T. tom Dieck and T. Petrie, *Geometric modules over the Burnside ring*, Invent. Math. 47 (1978), 273–287.
17. G. B. Segal, *Equivariant stable homotopy theory*, Actes du Congrès international des mathématiciens 1970, Nice, Tome 2, pp. 59–63, Gauthier–Villars, Paris, 1971.
18. G. B. Segal, *Categories and cohomology theories*, Topology 13 (1974), 293–312.
19. G. B. Segal, *Operations in stable homotopy theory*, in *New Developments in Topology*, London Math. Soc. Lecture Notes Series 11, pp. 105–110, Cambridge University Press, Cambridge, 1974.
20. C. Siebenneicher, *λ -Ringstrukturen auf dem Burnside ring der Permutationsdarstellungen einer endlichen Gruppe*, Math. Z. 146 (1976), 223–238.

21. D. Sullivan, *Genetics of homotopy theory and the Adams conjecture*, Ann. of Math. 100 (1974), 1–80.
22. O. Zariski and P. Samuel, *Commutative algebra II*, Graduate Texts in Mathematics 29, Springer-Verlag, New York - Heidelberg - Berlin, 1960.

ERKKI LAITINEN
UNIVERSITY OF HELSINKI
DEPARTMENT OF MATHEMATICS
HALLITUSKATU 15
SF-00100 HELSINKI 10
FINLAND